

apud

Reverse-genetic technologies utilizing gene knockout and knockdown have played a critical role in uncovering gene functions in many model and non-model systems. In *C. gigas*, most of the genetic studies have been carried out using the RNA interference (RNAi) technology to investigate the genetic regulation of germ cell development, reproduction, and salinity tolerance (Fabioux et al. [2009](#); Huvet et al. [2015](#); Zhao et al. [2017](#)). However, there are a number of caveats




Fig. 1 Schematic diagram of the MSTN-sgRNA and Twist-sgRNA target sites and sequences. The PAM sequences were shown in red, and the sgRNA target sequences were shown in blue. **a** MSTN-sgRNA was

targeted to the exon 3 (the antisense strand). **b** Twist-sgRNA was targeted to the exon 2 (the sense strand)

In Vitro Evaluation of sgRNA Efficacy

The gene editing efficacy of the sgRNA/Cas9 was tested in vitro using the Guide-it™ sgRNA Screening Kit (Clontech) according to the manufacturer’s instructions. Briefly, a 600–900 bp DNA fragment containing the sgRNA target sequence was amplified by PCR with the specific primers (Table 1). PCR products were purified using a SanPrep Column PCR Product Purification Kit (Sangon Biotech), and 100–250 ng amplicons were digested with sgRNA (50 ng) and Cas9 nuclease (250 ng) in a 15 µl reaction

volume at 37 °C for 1 h. The reaction was terminated by incubating at 80 °C for 5 min and stored at 4 °C for later use. Analysis of the enzymatic digestion was performed by electrophoresis on a 1.5% agarose gel.

Microinjection of sgRNA/Cas9 into Oyster Embryos

The sgRNA/Cas9 was delivered into oyster embryos by microinjection. An Olympus inverted microscope IX73 equipped with two micromanipulators for holding and microinjection pipettes (Narishige) was used. Holding and injection

Table 1 The primers used in this study

Primer name	Sequence (5'-3')
MSTN-sgRNA-F	<u>GATCACTAATACGACTCACTATAGGTTGTGAGCTGTAATCTAGGTTT</u> TAG AGCTAGAAAT
Twist-sgRNA-R	<u>GATCACTAATACGACTCACTATAGGACGATAACACGCACCTGTGTTT</u> TAG AGCTAGAAAT
sgRNA-common-R	AAAAGCACCGACTCGGTGCC
MSTN-sequencing-F	AACCCAACCTAGATTGCGGAC
MSTN-sequencing-R	TACGCACCAACGTTGAAGGTGA
Twist-sequencing-F	GACGGAATCCTGTCTTCTTGG
Twist-sequencing-R	TAGGAAAGAAACCTGACCATTTC
MSTN-sgRNA-verify-F	GACGAGATGCATGTTTCAATCGAT
MSTN-sgRNA-verify -R	TGCTGTAATCTTTCATAGGATGGAAC
Twist-sgRNA-verify-F	GAAACTTCAACACCCTTCACCT
Twist-sgRNA-verify-R	CAAACATACAAGGAGCAAGAATAGG

The underlined sequences in MSTN-sgRNA-F and Twist-sgRNA-R are 6-nt protective base. The red sequence in MSTN-sgRNA-F and Twist-sgRNA-R are the gene-specific target sequences within the respective sgRNAs

pipettes were prepared from Narishige glass capillaries with filament (GD-1, Narishige) using a Sutter P-97 micropipette puller (Sutter). The holding pipettes were further polished with a MR-900 microforge (Narishige), and the injection pipettes were further processed on an EG-401 microgrinder (Narishige). Microinjection was carried out using a PLI-100A Pico-Injector microinjector (Warner Instruments).

Alexa Fluor 488 phalloidin (AAT Bioquest) was injected at a concentration of 33 µg/µl. Approximately 0.1 nl of the fluorescent dye was microinjected in each oyster embryo. The sgRNAs and Cas9 protein (Guide-it™ Recombinant Cas9, Clontech) were mixed in equal volume and diluted in the injection buffer (20 mM HEPES, 150 mM KCl, and 0.5% phenol red) and then incubated for 15 min at 37 °C before use. Various concentrations of mstn-sgRNA and Twist-sgRNA were tested (Table 2). Different final concentrations of sgRNA and Cas9 protein were co-injected into 1–2-cell stage oyster embryos (Table 2). Approximately 0.1 nl of the sgRNA/Cas9 protein mixture was microinjected in each oyster embryo. The injected embryos were cultured in filtered seawater at 22 °C.

DNA Isolation

The genomic DNA was isolated from trochophores at approximately 12 h after fertilization using a proteinase K digestion method modified from Cai et al. (2018). Briefly, 20 µl of lysis buffer (1.5 mM MgCl₂, 10 mM Tris-HCl pH 8, 50 mM KCl, 0.5 mg/ml gelatin) was added to each larva in a 200-µl tube. The larvae were incubated at 94 °C for 10 min and 4 °C for 10 min. 2.5 µl of proteinase K (20 mg/ml) was added into each sample, and the samples were digested at 55 °C for 1 h, denatured at 94 °C for 10 min, and cooled down to 4 °C. The DNA samples were centrifuged, and 1 µl of the supernatant was used for each PCR reaction.

The genomic region covering the target sites and their flanking sequences were amplified using the respective

gene-specific primers (Table 1). Amplicons were purified and sequenced directly using the respective MSTN-F or Twist-F primers (Sangon Biotech). In addition, the PCR amplicons were cloned into pEASY vector. Thirty clones were randomly selected for plasmid isolation and DNA sequencing using the M13-F primer.

Results

In Vitro Cas9-sgRNA RNP Assay

To evaluate the efficacy of MSTN-sgRNA and Twist-sgRNA in guiding Cas9-induced gene editing at their respective target sites, we first tested these sgRNAs in vitro prior to microinjection into oyster embryos. DNA fragments containing the MSTN-sgRNA or Twist-sgRNA target sequences were amplified by PCR. The PCR products were incubated with MSTN-sgRNA/Cas9 or Twist-sgRNA/Cas9 ribonucleoprotein (RNP) complexes. The DNA products of the enzymatic digestion were analyzed by agarose gel. The results showed that incubating MSTN-sgRNA/Cas9 or Twist-sgRNA/Cas9 with their respective targets resulted in the production of two DNA bands with the expected size. In contrast, no DNA cleavage was detected when the PCR fragments were incubated with either Cas9, MSTN-sgRNA, or Twist-sgRNA alone (Fig. 2). Together, these results indicate that both MSTN-sgRNA and Twist-sgRNA could effectively guide Cas9 protein to cut DNA at their respective target sites.

Microinjection of Cas9-sgRNA RNP Complex

To assess the delivery of sgRNA/Cas9 RNP complex into oyster embryos by microinjection, we first tested microinjection of a fluorescent dye (Alexa Fluor 488 phalloidin) into

Table 2 Summary of injections with different concentrations of Cas9 protein and sgRNAs

Cas9 protein concentration (ng/µl)	sgRNAs concentration (ng/µl)	Survival (%)		Editing efficiency (%)	
		MSTN	Twist	MSTN	Twist
750	0	9/45 (20%)	8/40 (20%)	0/9 (0%)	0/8 (0%)
750	1500	8/40 (20%)	10/50 (20%)	8/30 (26.7%)	6/33 (18.2%)
750	375	8/41 (19.5%)	8/39 (20.5%)	0/8 (0%)	0/8 (0%)
375	0	9/37 (24.3%)	8/33 (24.2%)	0/9 (0%)	0/8 (0%)
375	1500	13/48 (27.1%)	12/46 (26.1%)	0/13 (0%)	0/12 (0%)
187	0	39/49 (79.6%)	35/40 (87.5%)	0/16 (0%)	0/16 (0%)
187	1500	40/50 (80%)	38/50 (76%)	0/16 (0%)	0/16 (0%)
0	1500	36/45 (80%)	34/40 (85%)	0/16 (0%)	0/16 (0%)
0	0	50/54 (92.6%)	45/50 (90%)	0/10 (0%)	0/10 (0%)




Fig. 2


fertilized oyster embryos at 1–2 cell stages. 0.1 nl of Alexa Fluor 488 phalloidin was microinjected into each embryo. The injected embryos were allowed to develop for 1 h and examined under a fluorescence microscope. A fluorescence signal was clearly detected in the injected embryos, and conversely the non-injected embryos showed no fluorescence signal (Fig. 3). The results indicated that reagents could be successfully injected into Pacific oyster embryos.

To investigate whether Cas9 protein or sgRNA is deleterious to oyster embryos, we decided to test the effect of injecting sgRNA or Cas9 protein alone on embryonic development and survival. The MSTN-sgRNA or Twist-sgRNA sgRNA was diluted at the final concentration of 1500 ng/μl, and 0.1 nl of the sgRNA was microinjected into each embryo. The injected embryos showed a survival rate of 80–85% to trochophores stage (Table 2). In contrast, injection of Cas9 protein at 375 ng/μl and 750 ng/μl resulted in a high mortality rate. Approximately 80% of the injected embryos died at the trochophores stage (Table 2). Reducing the Cas9 concentration to 187 ng/μl significantly improved the embryo survival to 79.6–87.5% (Table 2). Collectively, these data indicate that sgRNA and Cas9 RNPs can be delivered into oyster embryos by microinjection. The oyster embryos could tolerate sgRNA whereas a high dose of Cas9 protein is deleterious to the embryos.

CRISPR/Cas9
Target: MSTN, Twist

To determine whether microinjection of sgRNA/Cas9 could induce genetic mutations at their target genes in oyster, MSTN-sgRNA or Twist-sgRNA was mixed with a varying amount of Cas9 protein and injected into oyster embryos. The genomic regions surrounding the CRISPR targets were amplified by PCR and sequenced directly to investigate the successful rate of genome editing. The results revealed a clear dose-dependent effect of sgRNAs/Cas9 injection. At low doses of sgRNAs (375 ng/μl) and Cas9 protein (187 ng/μl and 375 ng/μl), no genetic mutation was observed in either *myostatin* or *Twist* genes in the injected embryos. However, injection of sgRNA at a higher concentration (1500 ng/μl) with a ninefold molar excess over Cas9 protein (750 ng/μl) successfully induced mutations in the target genes (Fig. 4), suggesting Cas9 RNPs induced mutations in a dose-dependent manner. Interestingly, we noted that injection of Cas9 protein at a higher concentration (750 ng/μl) and sgRNAs at a low concentration (375 ng/μl) resulted in no detectable mutations, suggesting that the molar ratio of sgRNAs to Cas9 protein is critical for efficient editing using the CRISPR/Cas9 system.

Fig. 3 Microinjection of Phalloidin 488 into oyster embryos. Phalloidin 488 was injected into oyster embryos. The injected and un-injected embryos were monitored under a fluorescence microscopy. **a** Bright field microscopy picture of embryos, and the red triangle indicated the injected embryos. **b** Fluorescence microscopy pictures of the injected embryos



To determine the successful rate of gene editing in the sgRNA (1500 ng/μl)/Cas9 (750 ng/μl) injected embryos, 30 larvae from the MSTN-sgRNA/Cas9 and 33 larvae from the Twist-sgRNA/Cas9 injected groups were selected for PCR and sequencing. Of the larvae sequenced, 26.7% and 18.2% of the injected individuals contained mutations at the target sites in MSTN and Twist genes, respectively. We noted that a higher concentration of sgRNA/Cas9 injection increased the rate of mutated embryos. However, this was associated with a reduced embryonic viability (~20%, Table 2).

C a ac, a, CRISPR/Ca 9-I . a I u
M. a, u, l, c, E b


To characterize the indel mutations induced by CRISPR/Cas9-mediated genome editing, we cloned the PCR amplicons from the injected larvae that displayed mutations on the sequencing chromatograph. Thirty clones were randomly selected for sequencing. Sequencing analysis showed

eight MSTN clones and six Twist clones contained DNA mutations at the target sites. As expected, various types of indel mutations were detected in MSTN and Twist genes, respectively (Fig. 5). The CRISPR/Cas9-induced mutations in the two target genes were predominantly small indel mutations ranging in size from 1 to 24 bp. Some of these mutations are frame-shift mutations that can disrupt the protein translation, and thus expected to be null mutations.

Discussion

In this study, we have established an effective approach to deliver sgRNA/Cas9 RNP complexes into *C. gigas* embryos by microinjection. We demonstrated that injection of sgRNA/Cas9 RNPs induced a dose-dependent mutations at the target genes. To our knowledge, this is the first report on gene editing in the Pacific oyster using the CRISPR technology. This approach should provide a solid basis for future

Fig. 4 Injection of MSTN-sgRNA/Cas9 or Twist-sgRNA/Cas9 RNPs induced DNA mutations at the target sites. MSTN-sgRNA/Cas9 or Twist-sgRNA/Cas9 RNPs were injected into oyster embryos. Thirty and 33 individual embryos were randomly selected for DNA isolation and PCR amplification of the DNA fragment covering the targeted regions. Representative DNA sequences of a control and a Cas9 RNPs injected embryos. Scrambled sequences were observed at –3 position of the MSTN-sgRNA target site (a) or Twist-sgRNA target site (b). WT meant sequence analysis of control embryos






Fig. 5 Characterization of indel mutations in the injected embryos. PCR amplicons from the MSTN-sgRNA/Cas9 or Twist-sgRNA/Cas9 RNPs injected embryos were cloned into the pEASY vector. Thirty clones were randomly selected for DNA sequencing. Alignments of the DNA sequences surrounding the targeting sites revealed five clones with indel

mutations in MSTN (**a**) and 4 clones with mutations in Twist genes (**b**). The wild-type (WT) sequence was in the top line, and mutant sequences

CRISPR/Cas9 studies of gene function in oysters and other bivalves, and potentially as a powerful tool for genetic engineering to improve oyster traits for aquaculture.

D² c
b M c **RNA/Ca 9** **O** **E b**

There are three major methods to introduce the CRISPR/Cas9 system into cells or organisms: (1) introduction of Cas9 and sgRNA as an all-in-one DNA plasmid expression vector, (2) introduction of Cas9 and sgRNA as in vitro transcribed RNAs, and (3) introduction of Cas9 protein and in vitro transcribed sgRNA as sgRNA/Cas9 ribonucleotide protein complexes (Hiruta et al. 2018). The third method via the delivery of sgRNA/Cas9 RNPs was found to be a simple and effective approach in CRISPR/Cas9 mediated gene editing. Direct delivery of sgRNA/Cas9 RNPs has been reported to yield a high editing efficiency in several different organisms such as mouse, zebrafish, water flea, and fungi (Aida et al. 2015; Burger et al. 2016; Hiruta et al. 2018; Wang et al. 2018). In addition, direct delivery of sgRNA/Cas9 RNPs by microinjection has several advantages. First, there is no need to consider the promoter and codon usage. Second, Cas9 and sgRNA are able to form a stable ribonucleoprotein in vitro, and the pre-loaded sgRNAs are possibly protected from degradation (Burger et al. 2016). Third, using the in vitro assembled sgRNA/Cas9 RNPs alleviates the possibility of integration of genetic material to the host genome (Wang et al. 2018). Consistent with these studies, we successfully applied the sgRNA/Cas9 RNPs system in targeted gene editing in the Pacific oyster.

Various methods, such as microinjection, electroporation, and lipofection, have been utilized for delivery of CRISPR/Cas9 into different organisms. Each method has its own advantages and disadvantages. For example, electroporation allows the simultaneous delivery of CRISPR/Cas9 systems into

a larger number of embryos. However, this method can cause significant cell death, and may not work for some marine species which are not tolerant to isotonic solution with low ionic strength for electroporation (Liu et al. 2017; Momose and Concordet 2016). Lipofection is relatively safe and easy to prepare, but low delivery efficiency has hampered its application. Microinjection is the direct injection of CRISPR/Cas9 into embryos with a high reproducibility and with minimal reagents using a micropipette at a microscopic level. Because of its simplicity and accuracy, microinjection has been used mostly so far. Certainly, there are several disadvantages of microinjection. Microinjection requires a high level of sophistication and manual skills and is only suitable for a limited number of embryos (Momose and Concordet 2016).

Our studies here support the idea that microinjection is an effective approach to deliver CRISPR/Cas9 in both model and non-model animals. However, there are several challenges for microinjection in *C. gigas*. Although a large number of eggs can be produced, the small size (< 50 μm in diameter) and opacity of the oocytes make the introduction of sgRNA/Cas9 RNPs by microinjection technically difficult. The first key factor for microinjection is the immobilization of eggs. In zebrafish, embryos can be easily held in wedge-shaped troughs made with 1.5% agarose. However, it is not suitable for oyster eggs because they are too small to be arranged in wedge-shaped troughs. In sea urchin, zygotes can be adhered to microinjection dishes coated with protamine sulfate or poly-L Lysine (Stepicheva and Song 2014; Hirsinger et al. 2015). We were not successful with this approach in oysters (data not shown here). In our study, the oyster eggs were held in place with a holding glass pipette by gentle suction. The next challenge is the tip size of the injection needle. Due to the small size of oyster eggs, the diameter of the needle tip must be very small ($\sim 1 \mu\text{m}$), which makes it easily clogged by oocyte cytoplasm and viscous injection solution especially at

the high Cas9 concentration. Therefore, a sharp and beveled pipette tip is preferable. Although microinjection is feasible as a practical method to deliver sgRNA/Cas9 RNPs into oyster embryos, it is, however, time-consuming and technically demanding to produce mutant oysters on a large scale. High throughput methods still need to be explored for applying gene editing technologies to improve oysters for aquaculture.

B a a c B l u M i a R a a E b
S a

Our data indicated that higher concentration of sgRNA/Cas9 injection is a key factor in inducing efficient genetic mutations in *C. gigas* embryos. We found that the sgRNA and Cas9 concentrations used in oyster microinjection were tenfold higher than that used in zebrafish in order to produce mutations (Cai et al. 2018). This could be explained by the smaller volume of sgRNA/Cas9 injected into each *C. gigas* embryo. In zebrafish, a typical volume of microinjection is 2–3 nl per embryos. In contrast, only 0.1 nl could be injected into the *C. gigas* embryo. This could be related to the lower efficiency in gene editing compared with zebrafish. Although successful induction of genetic mutations has been routinely accomplished in mouse embryos via microinjection of sgRNA/Cas9 at a moderate concentration comparable with that used in zebrafish, it should be noted that this is done with pronuclear injection into mouse embryos. Direction injection of sgRNA/Cas9 into the pronucleus allows a high concentration of the reagents in the nucleus and easy access to the genomic DNA.

We noted that a higher concentration of sgRNA/Cas9 injection increased the rate of mutated embryos. We found that the molar ratio of sgRNAs to Cas9 protein is critical for efficient editing using the CRISPR/Cas9 system. This is consistent with a previous report that six-fold molar excess of sgRNA over Cas9 protein resulted in maximizing the on-target mutation frequencies when delivered into cultured human leukemia K562 cells (Kim et al. 2014). However, a higher concentration of shRNA/Cas9 injection is associated with a reduced embryonic viability. A similar observation has been reported in monkeys (Midic et al. 2017). One feasible solution is the deployment of multiple sgRNAs (Zuo et al. 2017). Therefore, in future studies, it will be important to consider the deployment of multiple sgRNAs targeting the same gene. In addition, more sgRNAs can be designed and tested, because different sgRNAs show different efficiencies.

T C a a c i c I M a O
E b

We found that the CRISPR/Cas9-induced mutations were predominantly small indel mutations in these two target genes, ranging in size from 1 to 24 bp. This is very similar to the

results obtained in other invertebrate species (Lin and Su 2016; Hiruta et al. 2018). Because several different types of indels were observed from a single injected larva, it suggested that the indel mutations might occur independently in different cells. Moreover, we showed that only 20–25% of the sequenced clones were found to be mutated, suggesting the existence of mosaicism, which is common in gene-edited animals using the CRISPR/Cas9 system in injected embryos (Mehravar et al. 2018). Mosaicism in gene editing is not a problem if mutants are identified in the F1, F2, and F3 generations and used for gene functional studies.

Author Contributions Conceived and designed the experiments: QL and SD. Performed the experiments: HL, RX, CY, and HY. Analyzed the data: HL, HY, and SD. Wrote and revised the paper: HY, QL, and SD.

Funding Information This research was supported by grants from National Natural Science Foundation of China (31672649), Shandong Province (2017LZGC009), the Fundamental Research Funds for the Central Universities (201762014), and Taishan Scholars Seed Project of Shandong.

C a c i E c a S a a

All animal studies were carried out according to the guideline for the Care and Use of Laboratory Animals of Ocean University of China. The protocol was approved by the Institutional Animal Care and Use Committee of Ocean University of China.

Competing Interest The authors declare that there is no conflict of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Aida T, Chiyo K, Usami T, Ishikubo H, Imahashi R, Wada Y, Tanaka KF, Sakuma T, Yamamoto T, Tanaka K (2015) Cloning-free CRISPR/Cas system facilitates functional cassette knock-in in mice. *Genome Biol* 16:87
- Bono JM, Olesnick EC, Matzkin LM (2015) Connecting genotypes, phenotypes and fitness: harnessing the power of CRISPR/Cas9 genome editing. *Mol Ecol* 24:3810–3822
- Burger A, Lindsay H, Felker A, Hess C, Anders C, Chiavacci E, Zaugg J, Weber LM, Catena R, Jinek M, Robinson MD, Mosimann C (2016) Maximizing mutagenesis with solubilized CRISPR-Cas9 ribonucleoprotein complexes. *Development* 143:2025–2037
- Cai M, Si Y, Zhang J, Tian Z, Du S (2018) Zebrafish embryonic slow muscle is a rapid system for genetic analysis of sarcomere organization by CRISPR/Cas9, but not NgAgo. *Mar Biotechnol* 20:168–181
- Doudna JA, Charpentier E (2014) The new frontier of genome engineering with CRISPR-Cas9. *Science* 346:1258096
- Fabioux C, Corporeau C, Quillien V, Favrel P, Huvet A (2009) In vivo RNA interference in oyster-vasa silencing inhibits germ cell development. *FEBS J* 276:2566–2573
- FAO (2018) The state of world fisheries and aquaculture 2016. Food and Agriculture Organization of the United Nations, Rome

- Hirsinger E, Carvalho JE, Chevalier C, Lutfalla G, Nicolas JF, Peyri ras N, Schubert M (2015) Expression of fluorescent proteins in *Branchiostoma lanceolatum* by mRNA injection into unfertilized oocytes. *J Vis Exp* 95:52042
- Hiruta C, Kakui K, Tollefsen KE, Iguchi T (2018) Targeted gene disruption by use of CRISPR/Cas9 ribonucleoprotein complexes in the water flea *Daphnia pulex*. *Genes Cells* 23:494–502
- Huang X, Huan P, Liu B (2017) A comparative proteomic analysis reveals important proteins for the fertilization and early embryonic development of the oyster *Crassostrea gigas*. *Proteomics* 17:1600251
- Huvet A, B guel JP, Cavaleiro NP, Thomas Y, Quillien V, Boudry P, Alunno-Bruscia M, Fabioux C (2015) Disruption of amylase genes by RNA interference affects reproduction in the Pacific oyster *Crassostrea gigas*. *J Exp Biol* 218:1740–1747
- Kim S, Kim D, Cho SW, Kim J, Kim JS (2014) Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome Res* 24:1012–1019
- Li Q, Wang Q, Liu S, Kong L (2011) Selection response and realized heritability for growth in three stocks of the Pacific oyster *Crassostrea gigas*. *Fish Sci* 77:643–648
- Lin CY, Su YH (2016) Genome editing in sea urchin embryos by using a CRISPR/Cas9 system. *Dev Biol* 409:420–428
- Liu C, Zhang L, Liu H, Cheng K (2017) Delivery strategies of the CRISPR-Cas9 gene-editing system for therapeutic applications. *J Control Release* 266:17–26
- Mehravar M, Shirazi A, Nazari M, Banan M (2018) Mosaicism in CRISPR/Cas9-mediated genome editing. *Dev Biol* 445:156–162
- Midic U, Hung PH, Vincent KA, Goheen B, Schupp PG, Chen DD, Bauer DE, VandeVoort CA, Latham KE (2017) Quantitative assessment of timing, efficiency, specificity and genetic mosaicism of CRISPR/Cas9-mediated gene editing of hemoglobin beta gene in rhesus monkey embryos. *Hum Mol Genet* 26:2678–2689
- Momose T, Concordet JP (2016) Diving into marine genomics with CRISPR/Cas9 systems. *Mar Genomics* 30:55–65
- Perry KJ, Henry JQ (2015) CRISPR/Cas9-mediated genome modification in the mollusc, *Crepidula fornicata*. *Genesis* 53:237–244
- Riviere G, He Y, Tecchio S, Crowell E, Gras M, Sourdain P, Guo X, Favrel P (2017) Dynamics of DNA methylomes underlie oyster development. *PLoS Genet* 13:e1006807
- Song K, Li Y, Huang B, Li L, Zhang G (2017) Genetic and evolutionary patterns of innate immune genes in the Pacific oyster *Crassostrea gigas*. *Dev Comp Immunol* 77:17–22
- Stepicheva NA, Song JL (2014) High throughput microinjections of sea urchin zygotes. *J Vis Exp* 83:e50841
- Taning CNT, Van Eynde B, Yu N, Ma S, Smaghe G (2017) CRISPR/Cas9 in insects: applications, best practices and biosafety concerns. *J Insect Physiol* 98:245–257
- Wallis, de Paiva JS, van Prooijen BC, Ysebaert T, Smaal AC (2015) The ecosystem engineer *Crassostrea gigas* affects tidal flat morphology beyond the boundary of their reef structures. *Estuar Coasts* 38:941–950
- Wang Q, Li Q, Kong L, Yu R (2012) Response to selection for fast growth in the second generation of Pacific oyster (*Crassostrea gigas*). *J Ocean Univ China* 11:413–418
- Wang Q, Cao R, Ning X, You L, Mu C, Wang C, Wei L, Cong M, Wu H, Zhao J (2016) Effects of ocean acidification on immune responses of the Pacific oyster *Crassostrea gigas*. *Fish Shellfish Immunol* 49:24–33
- Wang Q, Cobine PA, Coleman JJ (2018) Efficient genome editing in *Fusarium oxysporum* based on CRISPR/Cas9 ribonucleoprotein complexes. *Fungal Genet Biol* 117:21–29
- Yue C, Li Q, Yu H (2018) Gonad transcriptome analysis of the Pacific oyster *Crassostrea gigas* identifies potential genes regulating the sex determination and differentiation process. *Mar Biotechnol* 20:206–219
- Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, Yang P, Zhang L, Wang X, Qi H, Xiong Z, Que H, Xie Y, Holland PW, Paps J, Zhu Y, Wu F, Chen Y, Wang J, Peng C, Meng J, Yang L, Liu J, Wen B, Zhang N, Huang Z, Zhu Q, Feng Y, Mount A, Hedgecock D, Xu Z, Liu Y, Domazet-Lo o T, Du Y, Sun X, Zhang S, Liu B, Cheng P, Jiang X, Li J, Fan D, Wang W, Fu W, Wang T, Wang B, Zhang J, Peng Z, Li Y, Li N, Wang J, Chen M, He Y, Tan F, Song X, Zheng Q, Huang R, Yang H, Du X, Chen L, Yang M, Gaffney PM, Wang S, Luo L, She Z, Ming Y, Huang W, Zhang S, Huang B, Zhang Y, Qu T, Ni P, Miao G, Wang J, Wang Q, Steinberg CE, Wang H, Li N, Qian L, Zhang G, Li Y, Yang H, Liu X, Wang J, Yin Y, Wang J (2012) The oyster genome reveals stress adaptation and complexity of shell formation. *Nature* 490:49–54
- Zhao X, Yu H, Kong L, Liu S, Li Q (2016) High throughput sequencing of small RNAs transcriptomes in two *Crassostrea* oysters identifies microRNAs involved in osmotic stress response. *Sci Rep* 6:22687
- Zhao X, Li Q, Meng Q, Yue C, Xu C (2017) Identification and expression of cysteine sulfinatase decarboxylase, possible regulation of taurine biosynthesis in *Crassostrea gigas* in response to low salinity. *Sci Rep* 7:5505
- Zuo E, Cai YJ, Li K, Wei Y, Wang BA, Sun Y, Liu Z, Liu J, Hu X, Wei W, Huo X, Shi L, Tang C, Liang D, Wang Y, Nie YH, Zhang CC, Yao X, Wang X, Zhou C, Ying W, Wang Q, Chen RC, Shen Q, Xu GL, Li J, Sun Q, Xiong ZQ, Yang H (2017) One-step generation of complete gene knockout mice and monkeys by CRISPR/Cas9-mediated gene editing with multiple sgRNAs. *Cell Res* 27:933–945