



# Epigenetic variation of wild populations of the Pacific oyster *Crassostrea gigas* determined by methylation-sensitive amplified polymorphism analysis

Xin Zhang<sup>1</sup> · Qi Li<sup>1,2</sup> · Lingfeng Kong<sup>1</sup> · Hong Yu<sup>1</sup>

Received: 16 July 2017 / Accepted: 10 November 2017 / Published online: 5 December 2017  
© Japanese Society of Fisheries Science 2017

## Abstract

Epigenetic variation can provide insights into ecological and evolutionary processes of eukaryote well-being, and can also be useful information for the conservation and management of wildlife. DNA methylation, a major epigenetic mechanism, plays a central role in gene regulation and phenotype variation in eukaryote genomes. However, in natural invertebrate populations, the potential role of epigenetic variation in DNA methylation polymorphisms and the association between this and genetic profiles remains largely unknown. In this work, we estimated epigenetic diversity and differentiation of the Pacific oyster *Crassostrea gigas*, from seven natural Chinese and Korean populations using the fluorescent-labeled methylation-sensitive amplified polymorphism (F-MSAP) technique. In total, 636 MSAP loci revealed an unambiguous genome-wide methylation status (31.32–43.23% on average) as well as extensive epigenetic polymorphism (approximately 97%) at 5'-CCGG-3' sites. The unweighted pair group method with arithmetic mean and principal coordinates analysis separated the seven *C. gigas* populations remarkably well, reflecting great epigenetic differentiation among populations, and in particular for the Korean population. Variation was higher in epigenetic structure than in genetic structure. Both multivariate analysis and a significantly different population molecular structure suggested that epigenetic variation might be determined by the combined effects of geographical barriers as well as current features and founder characteristics. A Mantel test showed a moderate but significant correlation between epigenetic and genetic profiles, indicating epigenetic variation coupled with genetic distribution in natural, wild *C. gigas* populations. This study provides some information on the role of DNA methylation in adaptive epigenetic divergence in wild marine bivalve populations.

**Keywords** Natural populations · Epigenetic differentiation · Conservation · DNA methylation · Genetic regulation

## Introduction

Environmental certainty and uncertainty generally represent considerable selective forces on natural populations (Via et al. 1995; Jaenisch and Bird 2003). Organisms can employ a suite of regulatory strategies, and modify their phenotype,

to cope with environmental fluctuations (Ghalambor et al. 2007). Recent work has indicated that epigenetic processes play crucial roles in phenotypic variation, which can lead to remarkable ecological and evolutionary success in natural plants and animals (Grant-Downton and Dickinson 2005; Bonasio et al. 2010; Feng et al. 2010; Faulk and Dolinoy 2011). Epigenetic status can provide insights into the ecological and evolutionary processes of eukaryote well-being, and is also of relevance to the conservation and management of wildlife.

Theoretically, epigenetic regulation is stably inheritable for gene products in the maintenance of genome integrity rather than in the modification of nucleotide sequences (Jablonka and Lamb 1998; Morris 2001). DNA methylation, a major epigenetic mechanism, consists mostly

of 5'-cytosine methylation at CCGG sites (Klose and Bird 2006). Studies (Rando and Verstrepen 2007; Nicotra et al. 2015) have shown that DNA methylation responds to environmental change and signal integration in the eukaryote genome by a change in pattern, and plays a central role in gene regulation and phenotype variation. As a general rule, hypermethylation is routinely described as being correlated with the down-regulation of gene expression (Gonzalzo and Jones 1997; Rajasethupathy et al. 2012). The methylation-sensitive amplified polymorphism (MSAP) assay has proven to be a powerful integrator of genome-wide epigenetic status (Reyna-Lopez et al. 1997). Recent studies using MSAP have shown some meaningful results in methylation variation and epigenetic population structure at CCGG sites of natural plants (Ochogavía et al. 2009; Schulz et al. 2014; Avramidou et al. 2015; Venetsky et al. 2015), vertebrates (Morán and Pérez-Figueroa 2011; Liu et al. 2012) and invertebrates (Díaz-Freije et al. 2014; Jiang et al. 2016). There is increasing interest in the role of epigenetic processes in ecology and evolution. To date, there are few reports concerning genome-wide methylation shown by MSAP in wild bivalve populations.

Historically, the Pacific oyster *Crassostrea gigas* is of great importance as a representative domesticated bivalve species. *C. gigas* is distributed worldwide (Chaney and Gracey 2011) and is thus an important species in the global fishery and aquaculture industries (Ren et al. 2010). In the past few decades, wild *C. gigas* populations have faced considerable challenges from natural and anthropological stressors, and have shown unexpected mortality as a consequence (Aranishi 2006; Guo et al. 2015). As an eurythermal and euryhaline sessile filter-feeding species, *C. gigas* has a strong capability to cope with dynamic hostile environments, such as intertidal zones, and exposure to a variety of stressors (Miossec et al. 2009; Kawabe et al. 2010; Li et al. 2013). A species' capability to respond to environmental fluctuations can provide information on ecological and evolutionary processes. A growing number of reports exploring the genetic structure and differentiation of wild oyster populations have indicated that their genetic diversity is ecogeographically structured and might be partly determined by dispersal, gene flow, climatic factors and currents

(Li et al. 2015; Zhong et al. 2016). However, their epigenetic profiles have scarcely been identified.

In this study, we assessed the epigenetic biodiversity and differentiation within and among natural, wild *C. gigas* populations from seven geographically isolated regions, using the fluorescent-labeled methylation-sensitive amplified polymorphism (F-MSAP) assay. Our study aimed at exploring the amount and structure of cytosine methylation at CCGG sites in a genome-wide manner, and finding correlations by comparing epigenetic and genetic differentiation. The observed epigenetic variation in the wild Pacific oyster is of potential value for the study of invertebrates.

## Materials and methods

### Oyster material and DNA extraction


*C. gigas* from seven geographically isolated sites in eastern China and Incheon, Korea, were sampled under permit. Adult oysters from Zhoushan (ZS), Dalian (DL), Qingdao (QD), Weihai (WH), Weifang (WF), and Ganyu (GY) in China, and Incheon (KI) in Korea, were selected from 2013 to 2016 (Table 1; Fig. 1). Adductor tissue was harvested and frozen at  $-80^{\circ}\text{C}$  for DNA extraction. Genomic DNA was extracted from adductor tissue using a modified phenol–chloroform protocol (Li et al. 2006).

### F-MSAP analysis

As with MSAP, F-MSAP involves the use of two methylation-sensitive restriction enzymes, *HpaII* and *MspI*. *HpaII*, which is inactive when either of the two cytosines is fully methylated, cleaves hemi-methylated DNA, whereas *MspI* cleaves 5'-C5mCGG-3'. The isoschizomers *HpaII* and *MspI* both recognize the tetranucleotide sequence 5'-CCGG-3', but display different sensitivity to cytosine methylation (Shaked et al. 2001; Jiang et al. 2016). The level of DNA methylation for each individual can be measured based on the number of polymorphic bands between the *MspI* and *HpaII* MSAP reactions. In this study, we followed the protocol provided by Jiang et al. (2016) that was established for the analysis

**Table 1** Sampled populations of the Pacific oyster *Crassostrea gigas*

Population	Location	Sampling period	Geographical coordinates
ZS	Zhoushan, Zhejiang Province, China	2013	30.0°N, 122.3°E
DL	Dalian, Liaoning Province, China	October 2016	38.9°N, 121.4°E
QD	Qingdao, Shandong Province, China	November 2016	35.7°N, 120.4°E
WH	Weihai, Shandong Province, China	May 2016	36.4°N, 121.3°E
WF	Weifang, Shandong Province, China	May 2016	37.1°N, 119.5°E
GY	Ganyu, Jiangsu Province, China	August 2016	34.8°N, 119.2°E
KI	Incheon, Korea	2014	36.4°N, 126.6°E



**Fig. 1** Map of the sampling locations and ocean currents. ZS Zhoushan, Zhejiang Province, China; DL Dalian, Province, China; QD Qingdao, Shandong Province, China; WH Weihai, Shandong Province, China; WF Weifang, Shandong Province, China; GY Ganyu, Jiangsu Province, China; KI Inchon, Korea; KC Kuroshio Current; TC Tsushima Current; YSWC Yellow Sea Warm Current; YSCC Yellow Sea Coastal Current; CCC China Coastal Current

of genome-wide oyster DNA methylation. Briefly, 100 ng of genomic DNA was digested with *EcoRI* + *HpaII/MspI* in a 10- $\mu$ L volume containing 2 U *EcoRI* (Thermo), 2 U of either *HpaII* or *MspI* enzyme (Thermo), and 10 $\times$  Tango buffer for 4 h at 37 °C. The *EcoRI* sticky ends and one of the *HpaII/MspI* sticky ends were ligated to the DNA after digestion by adding to each final digestion mixture 5  $\mu$ L of a mix containing 5  $\mu$ M of *EcoRI* adapter, 50  $\mu$ M of *HpaII/MspI* adapter (Table 2), 1 U of T4 DNA ligase (Thermo) and 10 $\times$  T4 DNA ligase buffer in a final volume of 20  $\mu$ L. The mixture was then incubated at 16 °C overnight. The digested-ligated DNA was then diluted at 1:10

with TE<sub>0.1</sub> and used as the template for the pre-amplification reaction. The pre-amplification reaction cycling conditions were 20 cycles at 94 °C for 20 s, 56 °C for 30 s and 72 °C for 2 min, followed by 60 °C incubation for 30 min. After the pre-amplification, the polymerase chain reaction products were diluted 1:10 with TE<sub>0.1</sub>. Selective amplification was performed for 2 min of denaturing at 94 °C, then ten cycles of 20 s at 94 °C, 30 s at 66 °C, and 2 min at 72 °C, with a 1 °C decrease in the annealing temperature of each cycle, followed by 20 cycles of 20 s at 94 °C, 30 s at 56 °C, and 2 min 72 °C, with 60 °C incubation for 30 min. All adapters and primers are listed in Table 2. Eight primer combinations were employed during the selective amplification stage. *EcoRI* selective primers were labeled using a 6-carboxy-fluorescein reporter molecule. The whole experiment was repeated twice in order to retain only the fully reproducible MSAP bands for further processing.

**Data collection and statistical analysis**

The F-MSAP reactions were loaded simultaneously onto an ABI 3130 Genetic Analyzer (Applied Biosystems) and the DNA methylation status of the loci was subsequently obtained with GeneMapper version 4.0 software. To reduce the potential impact of size homoplasy, only unambiguous and intense bands, ranging in size from 50 to 500 base pairs, were scored. We further converted the MSAP data into a binary matrix (Cervera et al. 2002).

To elucidate epigenetic structure among different geographically isolated regions, the epigenetic variation was calculated by the Dice (1945) and Jaccard (1908) coefficient algorithms, respectively. A group-clustering analysis was performed using the unweighted pair group method with arithmetic mean (UPGMA). Furthermore, principal coordinate analysis (PCoA) was used to provide an indication of genome-wide variability and to summarize the data in a few synthetic variables. All the above analyses were conducted using the numerical taxonomy system of

**Table 2** Adapters and primers used for methylation-sensitive amplified polymorphism

Adapter and primer	<i>EcoRI</i> (5'–3')	<i>HpaII/MspI</i> (5'–3')
Adapter I	CTCGTAGACTGCGTACC	GACGATGAGTCTAGAA
Adapter II	AATTGGTACGCAGTCTAC	CGTTCTAGACTCATC
Pre-amplification primer	GACTGCGTACCAATTCA	GATGAGTCTAGAACGGT
Selective primer	GACTGCGTACCAATTCATC	GATGAGTCTAGAACGGTGC
	GACTGCGTACCAATTCACG	GATGAGTCTAGAACGGTGT
	GACTGCGTACCAATTCACG	GATGAGTCTAGAACGGTAC
	GACTGCGTACCAATTCAGG	GATGAGTCTAGAACGGTAT
	GACTGCGTACCAATTCACT	GATGAGTCTAGAACGGTAG
	GACTGCGTACCAATTCACT	GATGAGTCTAGAACGGTGT
	GACTGCGTACCAATTCAAG	GATGAGTCTAGAACGGTAC
	GACTGCGTACCAATTCACA	GATGAGTCTAGAACGGTAT

multivariate programs–PC (NTSYS-PC) software package, version 2.10e (Applied Biostatistics, USA), and were conducted with  $10^3$  replicates. A Mantel test implemented in the isolation by distance (IBD) Web service program (Jensen et al. 2005) was performed to test the IBD model by correlating geographic distance to epigenetic distance and genetic distance.

In addition, to further explore the contribution of epigenetic and genetic profiles to the structure of the *C. gigas* populations, MSAP loci were then classified as either methylation-susceptible loci (MSL) or non-methylated loci (NML), depending on whether the observed proportion of methylated states across all samples exceeded a user-defined error rate-based threshold (5% by default) (Bonin et al. 2007; Pérez-Figueroa 2013; Díaz-Freije et al. 2014). Those fragments showing polymorphism of the same state at least twice were used for subsequent analyses (Herrera and Bazaga 2010; Díaz-Freije et al. 2014). Using the R package msap (<http://msap.r-forge.r-project.org>) (Pérez-Figueroa 2013), MSAP data were partitioned into the two distinct methylation types [MSL (DNA methylation pattern) and NML (genetic structure)] to study the variation of CCGG-methylation states and determine the correlation among all seven populations. Shown in Table 3 are: (1) the total number of loci, (2) the percentage of polymorphic loci ( $P_{MSL}$  and  $P_{NML}$ ), and (3) the mean Shannon's information index ( $I_{MSL}$  and  $I_{NML}$ ), (4) the multilocus analysis of molecular variance (AMOVA;  $\phi_{ST}$ ). Additionally, gene flow among populations was calculated based on the MSL and NML by POPGENE 1.32.

Finally, the similarity between epigenetic and genetic profiles was calculated with the standardized Mantel coefficient using  $10^3$  permutations for significance (Mantel 1967), using the NTSYS-PC software package. According to Cervera et al. (2002), Dice similarity coefficients (Sneath and Sokal 1973) were compared independently from both epigenetic and genetic profiles in the NTSYS-PC software package. Additionally, we calculated pairwise Nei's unbiased measures of genetic identity and genetic distance (Nei 1978) of both profiles by POPGENE 1.32 (Yeh et al. 1999; Zhong et al. 2016) and calculated pairwise AMOVAs. We then compared these data in the NTSYS-PC.

## Results

### Genome-wide analysis of methylation profiles

The level of methylation for each individual can be measured by the number of sites with polymorphic bands between the *MspI* and *HpaII* MSAP reactions out of the total number of MSAP sites. Examples of radioactively labeled and fluorescently labeled MSAP patterns are shown in Fig. 2. To this end, 636 reproducible methylation sites were analyzed in all populations. Namely, for each site, only variations which had originated from cytosine methylation (polymorphism between the *MspI* and *HpaII* MSAP reactions) were considered.

The average level of methylation varied statistically between all seven populations (Fig. 3). Notably, the level

**Table 3** Collection sites of wild *C. gigas* populations, total methylation-susceptible loci (MSL) diversity and comparison with non-methylated loci (NML) diversity indexes

Population	MSL				NML			
	Sites	$P_{MSL}$ (%)	$I_{MSL}$	AMOVA	Sites	$P_{NML}$ (%)	$I_{NML}$	AMOVA
ZS	328	67.38	$0.5743 \pm 0.1059$		308	33.77	$0.5335 \pm 0.1242$	
DL	347	70.89	$0.5835 \pm 0.0993$		289	42.91	$0.5407 \pm 0.1269$	
QD	391	74.17	$0.5749 \pm 0.1147$		245	34.29	$0.5404 \pm 0.1393$	
WH	347	70.03	$0.5863 \pm 0.1055$		289	50.52	$0.5562 \pm 0.1244$	
WF	371	71.16	$0.5797 \pm 0.1062$		265	36.98	$0.5366 \pm 0.1326$	
GY	377	74.54	$0.5729 \pm 0.1099$		259	48.06	$0.5369 \pm 0.1309$	
KI	352	49.72	$0.5859 \pm 0.1011$		284	26.41	$0.5171 \pm 0.1338$	
Mean	359	68.25			277	38.99		
Total	395	96.71	$0.5972 \pm 0.1126$		241	91.29	$0.4452 \pm 0.1607$	
Among groups				17.12				6.361
Within groups				41.94				18.26
				$\phi_{ST} = 0.2899,$ $P < 0.0001$				$\phi_{ST} = 0.2584,$ $P < 0.0001$

Analysis of molecular variance (AMOVA) among groups and within groups


$P_{MSL}$  (%) Percentage of polymorphic MSL loci,  $I_{MSL}$  Shannon's information index based on MSL loci,  $P_{NML}$  (%) percentage of polymorphic NML bands,  $I_{NML}$  Shannon's information index based on NML loci; for other abbreviations, see Table 1

## Clustering analysis

A phylogenetic tree was built based on the methylation patterns of total 636 CCGG sites from F-MSAP for all seven populations (Fig. 4; sample number—KI 1–8, ZS 9–16, DL 17–24, QD 25–32, WH 33–40, WF 41–48, and GY 49–56), in which the phylogenetic tree significantd. 1 rg/GSs0 Tc0 as higher

of the total methylation in Incheon, Korea was higher than that of populations from other areas ( $P < 0.01$ ). The total methylation level ranged from 31.32% (GY) to 43.23% (KI), with a mean of 36.24% at the species level. The average level of full methylation was somewhat similar, except for WF and KI populations ( $P < 0.01$ ). For hemi-methylation, the level of ZS, QD and KI was higher than that of the rest of the populations ( $P < 0.05$ ). There was a greater difference between hemi-methylation as compared with full methylation across the populations.





**Fig. 5** Principal coordinate analysis (PCoA) by MSAP in natural *C. gigas* populations. PCoA analysis used the Dice similarity measure in the numerical taxonomy system of multivariate program-PC software for the MSAP analysis. PCoA produces an ordination based on a distance or dissimilarity matrix where similar groups are clustered on a three-dimensional plot. The index indicates the different populations

The  $\phi_{ST}$  values between KI and the other populations were up to  $> 0.33$  and those of NML were up to  $> 0.27$  (Table 4), indicating large differentiation and variation between KI and the other populations. To some degree, the MSL results are in agreement with those of the UPGMA and PCoA. Additionally, AMOVA analysis for both MSL and NML showed that the vast majority of variation existed within populations; however, the differentiation level among populations was relatively low (Table 3). There was relatively limited calculated gene flow among the populations (MSL  $N_m = 1.6837$ , NML  $N_m = 0.9196$ ). IBD analyses showed that epigenetic and genetic distances were not significantly correlated with geographic distances ( $P_{epi} = 0.9370$ ,  $r_{epi} = 0.5165$ ;  $P = 0.8780$ ,  $r = 0.5896$ ), indicating that there were no significant phylogeographic structures of *C. gigas* in these areas. AMOVA for each population and among all populations revealed a higher variation in MSL than in NML. The Shannon index showed results similar to those in Table 3. The total Shannon index for methylation-susceptible polymorphic loci (MSL)

( $I_{epi} = 0.5971$ ) was statistically higher than the corresponding value for NML ( $I = 0.4452$ ) (Wilcoxon rank sum test with continuity correction  $W = 37283$ ,  $df = 6$ ,  $P < 0.0001$ ). The among-population Shannon index for MSL showed more abundance than that of the within-population index; the results for NML were opposite (Table 3).

### Correction between epigenetic and genetic profiles

The Mantel test was used to estimate the contribution of both epigenetic and genetic structure to natural *C. gigas* populations. Due to the deletion of MSL loci, we could not see any association between MSL and NML, as described above. So we calculated a similarity matrix using the transforming binary matrix described by Cervera et al. (2002) and the NML matrix as primitive data; the correlation between epigenetic and genetic profiles was evaluated based on Dice coefficients. Finally, the Mantel test showed some association between the transforming matrix and MSL profiles ( $r = 0.4234$ ,  $P = 0.0002$ ). A moderate but significant correlation was revealed between epigenetic and nucleotide sequence variation.

### Discussion

The genome-wide methylation of a total of 636 CCGG sites of 56 wild *C. gigas* individuals from seven geographically isolated populations was analyzed. Based on an unbiased F-MSAP analysis, the cytosine methylation variation was extensive and population specific, indicating a high level of epigenetic variation in all populations. Compared with domesticated *C. gigas* (~ 26% methylation in mass selection populations) (Jiang et al. 2013), the DNA methylation frequency of natural populations was relatively high, suggesting that *C. gigas* populations in natural environments might exhibit more epigenetic diversity. Similar results in other species (Shaked et al. 2001; He et al. 2015; Venetsky et al. 2015) are in agreement with ours. More differences were shown in hemi-methylation patterns than in fully methylated patterns, perhaps because isoschizomer *MspI* is insensitive to the methylation status of the internal cytosine

**Table 4** Pairwise  $\phi_{ST}$  values of MSL (below diagonal) and NML (above diagonal) among seven natural populations of Pacific oyster; for abbreviations, see Table 1

	ZS	DL	QD	WH	WF	GY	KI
ZS	*	0.2256	0.3373	0.2806	0.3664	0.3387	0.2852
DL	0.2896	*	0.2659	0.1539	0.2233	0.2469	0.2704
QD	0.2488	0.2328	*	0.2493	0.2883	0.2549	0.3468
WH	0.2252	0.1931	0.1991	*	0.2058	0.1982	0.275
WF	0.1892	0.175	0.1485	0.1928	*	0.2360	0.2924
GY	0.2586	0.2111	0.1832	0.1965	0.1357	*	0.3138
KI	0.4018	0.4230	0.3355	0.3486	0.3508	0.4069	*

(Blouin et al. 2010; Liu et al. 2012). The UPGMA and PCoA analyses showed significant clustering for most individuals. Considerable epigenetic differences between natural populations have been reported in previous studies, such as for wild female great roundleaf bat (Liu et al. 2012), wild barley (Li et al. 2008), etc. The specific methylation patterns might have been the result of a founder effect in different geographical regions, and might have an evolutionary role in the adaptation of natural populations.

Due to the lack of effective geographical barriers in the sea, geographic distance has been suggested to have an effect on molecular differentiation in marine shellfish populations (Palumbi 1994). Due to their geographical location, individuals of each *C. gigas* population could have unique patterns of inherited cytosine methylation through adaptation. Isolation by geographical distance is considered the most likely hypothesis to explain the differentiation among populations of different localities on the basis of a significant variation between molecular mechanisms and geographical distance (Arias et al. 2010). Analysis of the methylation status and epigenetic structure showed that KI was clearly separate from the Chinese oyster populations, indicating that geographical isolation might play a role in population epigenetic differentiation. However, no significant correlation was found between the epigenetic profile and geographic distance, suggesting a weak pattern of isolation by distance.

The inconsistency between specific epigenetic differentiation and geographical distance might indicate the influence of oceanographic and/or other environmental factors (Liu et al. 2012; Venetsky et al. 2015) which may have limited the degree of epigenetic divergence of *C. gigas*. GY is located at the junction of Jiangsu and Shandong Provinces, and the mesoscale oceanic eddy and vast area of coastal soil around this particular island could block gene flow among populations. This has also been reported for the genetic differentiation of populations of *Macra chinensis* (Ni et al. 2011) and *Octopus minor* (Gao et al. 2016). WF is located in the Bohai Sea, a semi-enclosed bay with a mildly sloping mud beach and shallow water, which means it is likely to have less water exchange and capacity for self-purification. Therefore, while the geographic distance from GY to WF is not small, the epigenetic variation between these locations is the lowest. On the other hand, there is relatively low epigenetic differentiation between WF and the other populations. As sessile organisms thriving in coastal oceans, the oysters may adapt to extensive and highly stressful environmental fluctuations (Miossec et al. 2009). Physical or biological factors may limit migration or free movement in benthic sessile organisms, and increase population differentiation of adult *C. gigas*. In this study, QD and WF showed more variation than the other populations. We inferred that other factors (sampling areas and human activities, etc.) could contribute to their extensive differentiation. Similar results have been

shown in previous studies (Li et al. 2015; Venetsky et al. 2015), and suggest that sampling area might be important for epigenetic population variation. The above suggested that geographic isolation could not, in isolation, explain epigenetic heterogeneity. In conclusion, it appears that there is no single factor that can explain *C. gigas* epigenetic variation, which is why the interactions between geographical isolation, oceanographic and life history characteristics have been the focus of this study.

It is interesting that epigenetic variation was structured into distinct among- and within-population components in *C. gigas*. This result was consistent with studies on plants (Li et al. 2008) and mammals (Liu et al. 2012). Previous studies have shown that epigenetic variance may not only be independent of (Cervera et al. 2002; Vaughn et al. 2007; Avramidou et al. 2015), but also dependent on, genetic variation (Li et al. 2008; Herrera and Bazaga 2010; Jiang et al. 2013; Schulz et al. 2014; Liu et al. 2015) in natural and cultivated populations. According to Richards (2006), there are two possible extreme cases for the interaction between epigenetic and genetic profiles: epigenetic variation can be strongly associated with genetic variation, or be completely uncoupled from genetic variation. Changes in epigenetic phenomena might occur much more frequently than in genetic factors, therefore, any changes in methylation within species might shed light on how they evolved (Liu et al. 2012; Avramidou et al. 2015; Venetsky et al. 2015). In the present study, e/GS1 gs-.014 Tw(201p Tc-.015 T ho)..al.



of drift could make epigenetic variations develop in parallel with genetic differences (Liu et al. 2012; Richards 2006) and lead to a moderate but significant correlation without a functional link between the two profiles. Besides the fact that both the epigenetic and genetic information was extracted from the same MSAP data set, different transformation probably leads to some biased estimates. Generally, our study supports the thesis that epigenetic variation is coupled with genetic variation. This study gives a first insight into the partitioning of epigenetic diversity in natural, wild Pacific oyster populations.

In conclusion, investigating DNA methylation polymorphisms in natural populations and evaluating the relevance of their epigenetic and genetic profiles are critical steps for assessing the potential role of epigenetic variation in natural invertebrate evolution. Using F-MSAP and multivariate analysis, we found that extensive epigenetic polymorphisms and distinct epigenetic structures might be partly determined by combined effects. Additionally, a moderate but significant association between epigenetic and genetic structure was verified in these natural oyster populations. This study provides some useful information on the role of DNA methylation in the adaptive epigenetic divergence of wild invertebrate populations.

**Acknowledgments** We thank the Marine Mollusk Resource Bank of Korea for providing oyster samples of Korea. This study was supported by grants from the National Natural Science Foundation of China (31772843), the Key Research and Development Program of Shandong Province (2016ZDJS06A06), the Industrial Development Project of Qingdao City (17-3-3-64-nsh), and the Fundamental Research Funds for the Central Universities (201762014).

## References

- Aranishi F (2006) A novel mitochondrial intergenic spacer reflecting population structure of Pacific oyster. *J Appl Genet* 47:119–123
- Arias A, Freire R, Méndez J, Insua A (2010) Isolation and characterization of microsatellite markers in the queen scallop *Aequipecten opercularis* and their application to a population genetic study. *Aquat Living Resour* 23:199–207
- Avramidou EV, Ganopoulos IV, Doulis AG, Tsaftaris AS, Aravanopoulos FA (2015) Beyond population genetics: natural epigenetic variation in wild cherry (*Prunus avium*). *Tree Genet Genomes* 11:95
- Blouin MS, Thuillier V, Cooper B, Amarasinghe V, Cluzel L, Araki H, Grunau C (2010) No evidence for large differences in genomic methylation between wild and hatchery steelhead (*Oncorhynchus mykiss*). *Can J Fish Aquat Sci* 67:217–224
- Bonasio R, Tu S, Reinberg D (2010) Molecular signals of epigenetic states. *Science* 330:612–616
- Bonin A, Ehrlich D, Manel S (2007) Statistical analysis of amplified fragment length polymorphism data: a toolbox for molecular ecologist and evolutionists. *Mol Ecol* 16:3737–3758
- Cervera MT, Ruiz-Garcia L, Martinez-Zapater J (2002) Analysis of DNA methylation in *Arabidopsis thaliana* based on methylation-sensitive AFLP markers. *Mol Genet Genom* 268:543–552
- Chaney ML, Gracey AY (2011) Mass mortality in Pacific oysters is associated with a specific gene expression signature. *Mol Ecol* 20:2942–2954
- Díaz-Freije E, Gestal C, Castellanos-Martínez S, Morán P (2014) The role of DNA methylation on *Octopus vulgaris* development and their perspectives. *Front Physiol* 5:62
- Dice LR (1945) Measures of the amount of ecologic association between species. *Ecology* 26:297–302
- Faulk C, Dolinoy DC (2011) Timing is everything: the when and how of environmentally induced changes in the epigenome of animals. *Epigenetics* 6:791–797
- Feng S, Jacobsen SE, Reik W (2010) Epigenetic reprogramming in plant and animal development. *Science* 330:622–627
- Gao XL, Zheng XD, Bo QK, Li Q (2016) Population genetics of the common long-armed octopus *Octopus minor* (Sasaki, 1920) (Cephalopoda: Octopoda) in Chinese waters based on microsatellite analysis. *Biochem Syst Ecol* 66:129–136
- Ghalambor CK, McKay JK, Carroll SP, Reznick DN (2007) Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Funct Ecol* 21:394–407
- Gonzalgo ML, Jones PA (1997) Mutagenic and epigenetic effects of DNA methylation. *Mutat Res* 386:107–118
- Grant-Downton R, Dickinson HG (2005) Epigenetics and its implications for plant biology. 1. The epigenetic network in plants. *Ann Bot* 96:1143–1164
- Guo XM, He Y, Zhang LL, Lelong C, Jouaux A (2015) Immune and stress responses in oysters with insights on adaptation. *Fish Shellfish Immun* 46:107–119
- He YY, Du Y, Li J, Liu P, Wang QY, Li ZX (2015) Analysis of DNA methylation in different tissues of *Fenneropenaeus chinensis* from the wild population and Huanghai no. 1. *Acta Oceanol Sin* 34:175–180
- Herrera CM, Bazaga P (2010) Epigenetic differentiation and relationship to adaptive genetic divergence in discrete populations of the violet *Viola cazorlensis*. *New Phytol* 187:867–876
- Jablonska E, Lamb MJ (1998) Epigenetic inheritance in evolution. *J Evol Biol* 11:159–183
- Jaccard P (1908) Nouvelles recherches sur la distribution florale. *Bull Soc Vaud Sci Nat* 44:223–270
- Jaenisch R, Bird A (2003) Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat Genet* 33:245–254
- Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. *BMC Genet* 6:13
- Jiang Q, Li Q, Yu H, Kong LF (2013) Genetic and epigenetic variation in mass selection populations of Pacific oyster *Crassostrea gigas*. *Genes Genom* 35:641–647
- Jiang Q, Li Q, Yu H, Kong LF (2016) Inheritance and variation of genomic DNA methylation in diploid and triploid Pacific oyster (*Crassostrea gigas*). *Mar Biotechnol* 18:124–132
- Kawabe S, Takada M, Shibuya R, Yokoyama Y (2010) Biochemical changes in oyster tissues and hemolymph during long-term air exposure. *Fish Sci* 76:841–855
- Klose RJ, Bird AP (2006) Genomic DNA methylation: the mark and its mediators. *Trends Biochem Sci* 31:89–97
- Li Q, Yu H, Yu RH (2006) Genetic variability assessed by microsatellites in cultured populations of the Pacific oyster (*Crassostrea gigas*) in China. *Aquaculture* 259:95–102
- Li YD, Shan XH, Liu XM, Hu LJ, Guo WL, Liu B (2008) Utility of the methylation-sensitive amplified polymorphism (MSAP) marker for detection of DNA methylation polymorphism and epigenetic population structure in a wild barley species (*Hordeum brevisubulatum*). *Ecol Res* 23:927–930
- Li Q, Zhao XL, Kong L, Yu H (2013) Transcriptomic response to stress in marine bivalves. *Invertebr Surviv J* 10:84–93

- Li S, Li Q, Yu H, Kong LF, Liu SK (2015) Genetic variation and population structure of the Pacific oyster *Crassostrea gigas* in the northwestern Pacific inferred from mitochondrial COI sequences. *Fish Sci* 81:1071–1082
- Liu S, Sun KP, Jiang TL, Ho JP, Liu B, Feng J (2012) Natural epigenetic variation in the female great roundleaf bat (*Hipposideros armiger*) populations. *Mol Genet Genom* 287:643–650
- Liu S, Sun KP, Jiang TL, Feng J (2015) Natural epigenetic variation in bats and its role in evolution. *J Exp Biol* 218:100–106
- Mantel N (1967) The detection of disease clustering and a generalized regression approach. *Cancer Res* 27:209–220
- Miossec L, Le Deuff RM, Goulletquer P (2009) Alien species alert: *Crassostrea gigas* (Pacific oyster). *ICES Coop Res Rep* 299:1017–6195
- Morán P, Pérez-Figueroa A (2011) Methylation changes associated with early maturation stages in the Atlantic salmon. *BMC Genet* 12:86
- Morris JR (2001) Genes, genetics, and epigenetics: a correspondence. *Science* 293:1103–1105
- Nei M (1978) Nei's unbiased measures of genetic identity and genetic distance. *Genetics* 89:583–590
- Ni LH, Li Q, Kong LF (2011) Microsatellites reveal fine-scale genetic structure of the Chinese surf clam *Macra chinensis* (Mollusca, Bivalvia, Macrtridae) in Northern China. *Mar Ecol* 32:488–497
- Nicotra AB, Segal DL, Hoyle GL, Schrey AW, Verhoeven KJ, Richards CL (2015) Adaptive plasticity and epigenetic variation in response to warming in an Alpine plant. *Ecol Evol* 5:634–647
- Ochogavía AC, Cervigni G, Selva JP, Echenique VC, Pessino SC (2009) Variation in cytosine methylation patterns during ploidy level conversions in *Eragrostis curvula*. *Plant Mol Biol* 70:17–29
- Palumbi SR (1994) Genetic divergence, reproductive isolation, and marine speciation. *Annu Rev Ecol Evol S* 25:547–572
- Pérez-Figueroa A (2013) msap: a tool for the statistical analysis of methylationsensitive amplified polymorphism data. *Mol Ecol Resour* 13:522–527
- Rajasethupathy P, Antonov I, Sheridan R, Frey S, Sander C, Tuschl T, Kandel ER (2012) A role for neuronal piRNAs in the epigenetic control of memory-related synaptic plasticity. *Cell* 149:693–707
- Rando OJ, Verstrepen KJ (2007) Timescales of genetic and epigenetic inheritance. *Cell* 128:655–668
- Ren JF, Liu X, Jiang F, Guo XM, Liu B (2010) Unusual conservation of mitochondrial gene order in *Crassostrea oysters*: evidence for recent speciation in Asia. *BMC Evol Biol* 10:394
- Reyna-Lopez G, Simpson J, Ruiz-Herrera J (1997) Differences in DNA methylation patterns are detectable during the dimorphic transition of fungi by amplification of restriction polymorphisms. *Mol Gen Genet* 253:703–710
- Richards EJ (2006) Opinion-inherited epigenetic variation-revisiting soft inheritance. *Nat Rev Genet* 7:395–401
- Rico L, Ogaya R, Barbeta A, Penuelas J (2014) Changes in DNA methylation fingerprint of *Quercus ilex* trees in response to experimental field drought simulating projected climate change. *Plant Biol* 16:419–427
- Schulz B, Eckstein RL, Durka W (2014) Epigenetic variation reflects dynamic habitat conditions in a rare floodplain herb. *Mol Ecol* 23:3523–3537
- Shaked H, Kashkush K, Ozkan H, Feldman M, Levy AA (2001) Sequence elimination and cytosine methylation are rapid and reproducible responses of the genome to wide hybridization and allopolyploidy in wheat. *Plant Cell* 13:1749–1759
- Sneath PHA, Sokal RR (1973) Numerical taxonomy. The principles and practice of numerical classification. Freeman, San Francisco
- Vaughn MW, Tanurdžić M, Lippman Z, Jiang H, Carrasquillo R, Rabonowicz PD, Dedhia N, McCombie WR, Agier N, Bulski A, Colot V, Doerge RW, Martienssen RA (2007) Epigenetic natural variation in *Arabidopsis thaliana*. *PLoS Biol* 5:e174
- Venetsky A, Levy-Zamir A, Khasdan V, Domb K, Kashkush K (2015) Structure and extent of DNA methylation-based epigenetic variation in wild emmer wheat (*T. turgidum* ssp. *dicoccoides*) populations. *BMC Plant Biol* 15:200
- Via S, Gomulkiewicz R, De Jong G, Scheiner SM, Schlichting CD, Van Tienderen PH (1995) Adaptive phenotypic plasticity: consensus and controversy. *Trends Ecol Evol* 10:212–217
- Wenzel MA, Piertney SB (2014) Fine-scale population epigenetic structure in relation to gastrointestinal parasite load in red grouse (*Lagopus lagopus scotica*). *Mol Ecol* 23:4256–4273
- Yeh FC, Yang RC, Boyle TBJ, Ye ZH, Mao JX (1999) POPGENE, version 1.32: the user-friendly software for population genetic analysis. Molecular Biology and Biotechnology Centre, University of Alberta, Canada
- Zhong XX, Feng DD, Yu H, Kong LF, Li Q (2016) Genetic variation and breeding signature in mass selection lines of the Pacific oyster (*Crassostrea gigas*) assessed by SNP markers. *PLoS One* 11:e0150868