

Effects of Salinity and Temperature on Growth and Survival of Juvenile Iwagaki Oyster *Crassostrea nippona*

WANG Tao¹⁾, and LI Qi^{1), 2), *}

1) Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, Qingdao 266003, China

2) Laboratory for Marine Fisheries Science and Food Production Processes, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266003, China

Abstract Iwagaki oyster *Crassostrea nippona* occurs naturally along the coasts of Japan and Korea. Because of its unique flavor, delicious taste, edibility during the summer and high commercial value, it has been identified as a potential aquaculture species. To determine the optimum aquaculture conditions and provide necessary information for mass production of the juvenile, the effects of six salinities (15, 20, 25, 30, 35 and 40) and five temperatures (16, 20, 24, 28 and 32°C) on growth and survival of juvenile *C. nippona* were examined in this study. In the salinity experiment, the largest values of mean shell height and growth rate were observed at salinity 25 (20.96 ± 0.36 mm and $172.0 \mu\text{m d}^{-1}$, respectively), which were significantly different ($P < 0.05$) with those of other treatments, except at salinity 30 (20.56 ± 1.05 mm and $160.3 \mu\text{m d}^{-1}$, respectively) ($P > 0.05$). The maximum survival rate 84.44% was always observed at salinity 20, and there was no significant difference ($P > 0.05$) in survival rate among salinities varying between 15 and 35. In the temperature-related experiments, the highest growth and survival rates of juvenile were observed at 24°C ($180.8 \mu\text{m d}^{-1}$ and 84.4%) and 28°C ($190.7 \mu\text{m d}^{-1}$ and 83.3%), respectively, on day 20, and showed significantly ($P < 0.05$) larger size and higher survival rate than any other groups. Both juvenile survival and growth were significantly depressed at extreme salinities (15, 40) and temperatures (16°C, 32°C). Based on the results of the present study, a salinity range from 25 to 30 and a temperature range from 24 to 28°C are considered optimal conditions for survival and growth of juvenile *C. nippona*.

Key words *Crassostrea nippona*; juvenile; salinity; temperature; survival; growth

1 Introduction

Iwagaki oyster *Crassostrea nippona* belongs to Bivalvia, Ostreidae, and is a large sessile oyster inhabiting intertidal hard grounds and reefs along the coast of East Asia including Japan and Korea (Itoh *et al.*, 2004). Because of its unique flavor, delicious taste and marketability in summer when Pacific oyster *C. gigas* is unavailable, the commercial price of *C. nippona* is estimated as high as five folds of *C. gigas* in Japan (Itoh *et al.*, 2004). Thus this species has a market prospect and high potential value of large scale farming. Traditionally, *C. nippona* farming is largely dependent on natural seeds (Tanaka *et al.*, 2010). However, such seed collection is labor intensive, often unreliable and available for a short season. Moreover, with the increasing interest of *C. nippona* culture, natural seed collection may not satisfy the demand of aquaculture, and seed production is always in short supply due to the deficiency of facilities (Tanaka *et al.*, 2010; Fujiwara, 1995). These issues have significantly

constrained the cultivation of *C. nippona*, hence, developing proper breeding, nursery, and aquaculture techniques for mass production of juveniles are crucial for meeting the success of *C. nippona* farming.

Autecological study of bivalves has clearly demonstrated that environmental factors play an important role in the development, growth and survival of aquatic animals (Tang *et al.*, 2012). Among them, salinity and temperature are considered to be the most important physical parameters affecting the physiological responses and survival of aquatic organisms, which have been described as 'master factors' (Re *et al.*, 2005; Kinne, 1964). Salinity imposes an additional metabolic load (Bao and You, 2004) and affects biological activities, including those related to immune responses (Taylor *et al.*, 2007), fertilization (Verween *et al.*, 2007), development of embryos (Davis and Calabrese, 1969), survival and growth of larvae and juvenile (Huo *et al.*, 2014). Temperature modifies energy flow, which regulates the rate of biological processes (Scheltema, 1967). Accordingly, the effects of these two factors have been described for numerous marine species (Ko *et al.*, 2014; Huo *et al.*, 2014; Xu *et al.*, 2011). For *C. nippona*, although previous studies have documented seasonal variations in reproductive activity and bio-

* Corresponding author. Tel: 0086-532-82031622
E-mail: qili66@ouc.edu.cn

chemical composition, karyotype, and culture method
(Itoh

different temperatures of 20°C to 32°C on day 5, 10 and 15, but a significant effect of temperature was observed after day 20. Among all treatments, it was noted that on day 20, survival rate was significantly ($P < 0.05$) higher at 24 and 28°C than that at 20 and 32°C, ranging from 83% to 84%. Survival rate of the juvenile was longer at 20°C than that at 32°C, whereas survival rate displayed no significant ($P > 0.05$) difference between these two treatments.

Table 3 Survival and growth rates of *Crassostrea nippona* juvenile held at different temperatures

| Temperature (°C) | Growth rate ($\mu\text{m d}^{-1}$) | Survival rate (%) | | | |
|------------------|--------------------------------------|---------------------|----------------------|----------------------|----------------------|
| | | 5 d | 10 d | 15 d | 20 d |
| 16 | 73.3±8.7 ^c | 86±5.1 ^a | 52±15.8 ^a | 50±15.3 ^a | 43±12.0 ^a |
| 20 | 124.3±18.7 ^b | 88±3.9 ^a | 81±10.2 ^b | 73±8.8 ^b | 70±5.8 ^b |
| 24 | 180.8±14.7 ^a | 91±5.1 ^a | 88±5.1 ^b | 87±6.7 ^b | 84±5.1 ^c |
| 28 | 190.7±18.6 ^a | 91±6.9 ^a | 89±6.9 ^b | 86±3.9 ^b | 83±3.3 ^c |
| 32 | 73.0±5.6 ^c | 92±1.9 ^a | 92±1.9 ^b | 76±1.9 ^b | 69±1.9 ^b |

Notes: Within a column, values followed by different letters are significantly different ($P < 0.05$, $n = 30$). Data are given as Mean ± SE.

4 Discussion

According to the observed mortalities and daily growth rates, different salinities have significant effects on juvenile *C. nippona*. Our findings agree with those of Yao *et al.* (2015) who studied the effect of salinity on larval growth, survival and development of the oysters *C. gigas* and *C. ariakensis*. Previous studies have demonstrated that oysters are generally euryhaline species, although different species may differ in the optimal salinity that they prefer (Xu *et al.*, 2011). For example, European flat oyster *Ostrea edulis* larvae had the high growth rates and survival at an optimum salinities of 24–33 (Walne, 1956). Sydney rock oyster *Saccostrea glomerata* larvae had the highest growth rates at salinities 23–39, and the highest survival rates at salinities 27–39 (Nell and Holliday, 1988). Mangrove oyster *C. rhizophorae* had the highest embryo development rates at salinities of 25–37 (Dos Santos and Nascimento, 1985). For *C. nippona*, there was no significant difference among different salinities of 15–35 after 20 days of cultivation, indicating that 15–35 is the optimum salinity for the survival of juvenile. However, the highest growth rates were observed at salinity of 25 and 30. When the salinity were adjusted to be lower or higher than this range, growth rate of juvenile was significantly affected, resulting in a decreased shell length. Therefore, a salinity range of 25–30 is the optimum salinity for the growth of juvenile. Similarly, in Kumamoto oyster *C. sikamea*, salinities higher than 25 had significant negative effects on the larvae development and growth (Xu *et al.*, 2011). The survival and growth of spat, juvenile and adult *C. virginica* were observed to be depressed directly with lower salinity. Combined with suboptimal elevated temperature, lower salinities have negative effects in all size classes of *C. virginica*, resulting in dramatically greater mortality (Rybovich *et al.*, 2016).

In the present study, rearing *C. nippona* in the lowest (15 and 20), medium (25 and 30), and the highest salinities (35 and 40) had significant effects on juvenile growth. A similar result was also observed in *C. nippona* larvae, in which the lowest growth rate and survival rate were measured at salinities of 14, 18 and 34 on day 13 (Wang *et al.*, unpublished data). The mechanisms through which salinity affects marine mollusks remain unclear. However, previous studies indicate that there were two possible reasons for decreased growth rates and increased mortality under extreme salinity conditions. Firstly, reduced feeding rates could be a major driver for reduced growth rates in extreme salinity since oysters completely seal themselves off by closing their shells when salinity drops too low or increases too high (Berger and Kharazova, 1997). For example, in the European flat oyster *Ostrea edulis*, salinity were found to significantly affect absorption efficiency, filtration rate, and metabolic response, further lead to retarder growth of larvae. (Hutchinson and Hawkins, 1992). Low energy utilization efficiency of the body could be the other reason. Like other marine molluscs, *C. nippona* was an osmotic conformer (Berger and Kharazova, 1997). Additional energy is required for the maintenance of water and mineral balance in body fluids and cells when juvenile *C. nippona* were cultivated at suboptimal salinity conditions (Berger and Kharazova, 1997). This conclusion agrees with that of Forcucci and Lawrence (1986), who unequivocally demonstrated that either additional energy is required for the maintenance or energy is used inefficiently that can cause the slow growth of *Luidia clathrata* when salinity was adjusted to a level that was lower or higher than optimal salinity ranges.

Temperature is another important environmental variable that can influence juvenile survival and growth. In the present study, minimum growth and survival rate were observed at 16°C, and as temperature increased, these values reached the maximum at 24 and 28°C. This agrees with many other studies on the growth of mollusks that juveniles generally grew more quickly at the higher temperatures (Klinzing and Pechenik, 2000; Pechenik and Heyman, 1987). Davis and Calabrese (1969) suggested that the failure of marine bivalve species to grow at low temperatures appears to be a result of their inability to digest the available food, although they can survive and ingest food for a long time. The appropriate improvement of cultivation temperature could promote the activities of certain digestive enzymes, and increase the assimilation efficiency of the organism, subsequently leading to an increase in somatic growth (Bayne *et al.*, 1999). However, with the temperature rising from 28 to 32°C, survival and growth of juvenile decreased sharply. This has been explained by the increase in growth rate (8.3(x)2r3.7(h)-1.7(e)6(i)85 1th susalir

to the protein synthesis, which resulted in the depletion of glycogen reserves and protein content in tissues, and eventually led to less favorable energy balance and smaller maximum size (Ivanina *et al.*, 2013). Therefore, the temperature of 24–28°C is optimum for the growth and survival of juvenile *C. nippona*. Similarly, optimum larval development and settlement of the oyster *C. gigas* occurred at 27°C (Rico-Villa *et al.*, 2009). For *O. edulis*, the average rate of growth of larvae increased progressively as the temperature increased from 10 to 25 or 27.5°C and then decreased at 30 and 32.5°C (Davis and Calabrese, 1969).

Another notable result in the present study is that juvenile cultivated at 16°C displayed lower survival rates than at 32°C, even though there was no significant difference between these two extreme temperatures on growth rates. The present result indicated that juvenile *C. nippona* had greater adaptability to high temperature than low temperature. This may be related to the summer breeding habits of *C. nippona*. The parental *C. nippona* has a sexual maturation between August and September, which is the period with the highest water temperature of the year (Okumura *et al.*, 2005). Similar phenomenon also shown in other marine bivalves, including *Saccostrea glomerata* and *Tegillarca granosa* (Parker *et al.*, 2009; You *et al.*, 2001).

In summary, the results of this study clearly demonstrate that juveniles of *C. nippona* can tolerate a wide range of salinities (15–35) and a wide range of temperatures (20–32°C), while exhibit larger shell height and better survival at salinities 25–30 and 24–28°C. Hence, culturing this species at lower salinities of 25–30 and higher temperatures of 24–28°C will likely produce a high yield and benefit juvenile *C. nippona*.

Acknowledgements

This study was supported by the grants from the Key Research and Development Program of Shandong Province (No. 2016ZDJS06A06), the National Natural Science Foundation of China (No. 31772843), and the Major Project for Tianjin Seed Technology (No. 15ZXZYNC0050).

References

- Adachi, K., Yokoi, T., Inoue, K., Yoshinaga, T., and Okumura, S. I., 2014. Karyotype revision in the Iwagaki oyster *Crassostrea nippona*. *Chromosome Science*, **17**: 9-10.
- Bao, Y. B., and You, Z. J., 2004. Influences of several environmental factors on growth in marine shellfish larvae. *Fisheries Science*, **23**: 39-41.
- Bayne, B. L., Svensson, S., and Nell, J. A., 1999. The physiological basis for faster growth in the Sydney rock oyster, *Saccostrea commercialis*. *The Biological Bulletin*, **197**: 377-387.
- Berger, V. J., and Kharazova, A. D., 1997. Mechanisms of salinity adaptations in marine molluscs. *Interactions and Adaptation Strategies of Marine Organisms*, **355**: 115-126.
- Davis, H. C., and Calabrese, A., 1969. Survival and growth of larvae of the European oyster (*Ostrea edulis* L.) at different temperatures. *The Biological Bulletin*, **136**: 193-199.
- Dos Santos, A. E., and Nascimento, I. A., 1985. Influence of gamete density, salinity and temperature on the normal embryonic development of the mangrove oyster *Crassostrea rhizophorae* Guilding, 1828. *Aquaculture*, **47**: 335-352.
- Forcucci, D., and Lawrence, J. M., 1986. Effect of low salinity on the activity, feeding, growth and absorption efficiency of *Luidia clathrata* (Echinodermata: Asteroidea). *Marine Biology*, **92**: 315-321.
- Fujiwara, M., 1995. The problems in seed production of Iwa oyster *Crassostrea nippona*. *Bulletin of the Kyoto Institute of Oceanic and Fishery Science*, **18**: 14-21 (in Japanese).
- Gruffydd, L. D., and Beaumont, A. R., 1972. A method for rearing *Pecten maximus* larvae in the laboratory. *Marine Biology*, **15**: 350-355.
- Huo, Z. M., Wang, Z. P., Liang, J., Zhang, Y. H., Shen, J. P., Yao, T., Su, J. Q., and Yu, R. H., 2014. Effects of salinity on embryonic development, survival, and growth of *Crassostrea hongkongensis*. *Journal of Ocean University of China*, **13**: 666-670.
- Hutchinson, S., and Hawkins, L. E., 1992. Quantification of the physiological responses of the European flat oyster *Ostrea edulis* L. to temperature and salinity. *Journal of Molluscan Studies*, **58**: 215-226.
- Itoh, N., Tun, K. L., Komiyama, H., Ueki, N., and Ogawa, K., 2004. An ovarian infection in the Iwagaki oyster, *Crassostrea nippona*, with the protozoan parasite *Marteilioides chungmuensis*. *Journal of Fish Diseases*, **27**: 311-314.
- Ivanina, A. V., Dickinson, G. H., Matoo, O. B., Bagwe, R., Dickinson, A., Beniash, E., and Sokolova, I. M., 2013. Interactive effects of elevated temperature and CO₂ levels on energy metabolism and biomineralization of marine bivalves *Crassostrea virginica* and *Mercenaria*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, **166**: 101-111.
- Kinne, O., 1964. The effects of temperature and salinity on marine and brackish water animals: II. Salinity and temperature-salinity combinations. *Oceanography Marine Biology Annual Review*, **2**: 281-339.
- Klinzing, M. S. E., and Pechenik, J. A., 2000. Evaluating whether velar lobe size indicates food limitation among larvae of the marine gastropod *Crepidula fornicata*. *Journal of Experimental Marine Biology and Ecology*, **252**: 255-279.
- Ko, G. W., Dineshram, R., Campanati, C., Chan, V. B., Havenhand, J., and Thiyagarajan, V., 2014. Interactive effects of ocean acidification, elevated temperature, and reduced salinity on early-life stages of the pacific oyster. *Environmental Science and Technology*, **48**: 10079-10088.
- Li, W. J., 2007. Biology and cultivation of oyster *Crassostrea nippona*. *Fisheries Science*, **12**: 689-690 (in Chinese).
- Nell, J. A., and Holliday, J. E., 1988. Effects of salinity on the growth and survival of Sydney rock oyster *Saccostrea commercialis* and Pacific oyster *Crassostrea gigas* larvae and spat. *Aquaculture*, **68**: 39-44.
- Okumura, T., Miura, N., Semura, H., and Kishimoto, O., 2005. Seasonal changes in glycogen contents of the Iwagaki oyster, *Crassostrea nippona* from the coast of Toga Bay, Konoura, Tomari, and Oki Islands in the Sea of Japan. *Nippon Suisan Gakkaishi*, **71**: 363-368 (in Japanese).
- Parker, L. M., Ross, P. M., and O'Connor, W. A., 2009. The effect of ocean acidification and temperature on the fertilization and embryonic development of the Sydney rock oyster *Saccostrea glomerata* (Gould 1850). *Global Change Biology*,

- 15: 2123-2136.
- Pechenik, J. A., and Heyman, W. D., 1987. Using KCI to determine size at competence for larvae the marine gastropod *Crepidula fornicata* (L.). *Journal of Experimental Marine Biology and Ecology*, **112**: 27-38.
- Re, A. D., Diaz, F., Sierra, E., Rodríguez, J., and Perez, E., 2005. Effect of salinity and temperature on thermal tolerance of brown shrimp *Farfantepenaeus aztecus* (Ives) (Crustacea, Penaeidae). *Journal of Thermal Biology*, **30**: 618-622.
- Rico-Villa, B., Pouvreau, S., and Robert, R., 2009. Influence of food density and temperature on ingestion, growth and settlement of Pacific oyster larvae, *Crassostrea gigas*. *Aquaculture*, **287**: 395-401.
- Rybovich, M., Peyre, M. K. L., Hall, S. G., and Peyre, J. F. L., 2016. Increased temperatures combined with lowered salinities differentially impact oyster size class growth and mortality. *Journal of Shellfish Research*, **35**: 101-113.
- Scheltema, R. S., 1967. The relationship of temperature to the larval development of *Nassarius obsoletus* (Gastropoda). *The Biological Bulletin*, **132**: 253-265.
- Tanaka, M., Hisada, T., and Fujiwara, M., 2010. Possibility of natural spat collection of 'Iwagaki' oyster *Crassostrea nippona* in western Wakasa Bay. *Bulletin of the Kyoto Institute of Oceanic and Fishery Science*, **32**: 17-22 (in Japanese).
- Tang, B. J., Yin, F., Gui, C. S., Chen, X. Z., and Liu, B. Z., 2012. Hatching, growth and hatchling dietary preference in *Hemifusus ternatanus* (Gmelin, 1791). *Aquaculture*, **326**: 141-147.
- Taylor, J. F., Needham, M. P., North, B. P., Morgan, A., Thompson, K., and Migaud, H., 2007. The influence of ploidy on saltwater adaptation, acute stress response and immune function following seawater transfer in non-smolting rainbow trout. *General and Comparative Endocrinology*, **152**: 314-325.
- Verween, A., Vincx, M., and Degraer, S., 2007. The effect of temperature and salinity on the survival of *Mytilopsis leucophaeata* larvae (Mollusca, Bivalvia): The search for environmental limits. *Journal of Experimental Marine Biology and Ecology*, **348**: 111-120.
- Walne, P. R., 1956. Experimental rearing of the larvae of *Ostrea edulis* in the laboratory. *Fisheries Investigation Series II*, **9**: 23.
- Xu, F., Guo, X. M., Li, L., and Zhang, G. F., 2011. Effects of salinity on larvae of the oysters *Crassostrea ariakensis*, *C. sikamea* and the hybrid cross. *Marine Biology Research*, **7**: 796-803.
- Yao, T., Wang, Z. P., Yan, X. W., Li, D. C., Zhang, Y. H., Huo, Z. M., Su, J. Q., and Yu, R. H., 2015. Effect of salinity on growth and survival of *Crassostrea gigas*, *C. ariakensis* and juvenile hybrids. *Acta Ecologica Sinica*, **5**: 1581-1586 (in Chinese).

(vl of