# Genome-Wide Scan for Positively Selected Genes in Sessile Molluscs Highlights the Genetic Basis for Their Adaptation to Attached Lifestyle

YU Hong<sup>1), 2)</sup>, and LI Qi<sup>1), 2), \*</sup>

1) Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, Qingdao 266003, China

2) Laboratory for Marine Fisheries Science and Food Production Processes, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266003, China

(Received August 3, 2017; revised November 13, 2017; accepted April 10, 2018) © Ocean University of China, Science Press and Springer-Verlag GmbH Germany 2018

Abstract Molluscs are one of the most diverse groups of animals and exhibit a rich and diverse variety of form and lifestyle. Most molluscs live with a free-moving lifestyle, while some molluscs are sessile. The adaptation to the two distinct lifestyles required complex changes, from molecules to organs, and physiology to morphology. In this study, we conducted a genome-wide scan for positive selection by comparing the available genomes of two sessile molluscs with four free-moving molluscs. A total of 40 genes were identified undergoing positive selection in the sessile molluscs by the branch-site model. Functional characterization showed that they were mainly enriched in two pathways, Oxidative phosphorylation (OXPHOS) and TGF-beta signaling pathway. The unexpected positive selection on OXPHOS genes in sessile molluscs suggests that the adaptation of OXPHOS involves many factors be yond enhancing ATP production. A modified OXPHOS regulatory system may allow sessile molluscs to better cope with biotic and abiotic stresses. Moreover, positively selected genes in TGF-beta signaling pathway probably have played a key role in the patterning of body plans and growth in metazoans. We speculate that these genes are associated with the body structure and organic adaptation to a sedentary lifestyle in sessile bivalve molluscs.

Key words molluscs; adaptation evolution; positive selection; sessile lifestyle

# 1 Introduction

Mollusca is the second largest phylum in the animal kingdom, and one of the most diverse groups of animals. Molluscs exhibit a rich and diverse variety of form and lifestyle. The molluscan body plan is enormously varied, and extremely plastic and adaptable, which has led to their remarkable evolutionary success, representing about 200000 living species (Ponder and Lindberg, 2008). Based on the moving ability, molluscs can be categorized into free-moving and sessile forms. In general, most of the molluscs are free-moving (*e.g.*, squids, cuttlefishes, snails, and sea slugs), while some moaet al., 2012). The adaptations can

lecules to organs. In molluscs, most sessile ar feeding with hypertrophied gills (Ponder 2008), and their head and foot have degen-

> \* Corresponding author. Tel: 0086-532-82031622 E-mail: gili66@ouc.edu.cn

erated. Changes in energy metabolism have inevitably occurred between the sessile and free-moving molluscs.

nal to diurnal circadian rhythm in mammals, seasonal and non-seasonal reproductive behaviors (Shen et al., 2012; Sun et al., 2013; Nery et al., 2013; Cao et al., 2015; Meng et al., 2015). However, our current understanding of selective imprints is mainly limited to mammals. The molluscs are one of the most diverse phyla on Earth and play an important role as model organisms in scientific research (Ponder and Lindberg, 2008), but their genetic adaptation to ecological niches is largely unknown. Recently, emerging high-throughput sequencing technology and bioinformatics tools have undoubtedly facilitated the study on the genetic basis of how molluscs adapt to diverse lifestyles. To date, six molluscan genomes (the Pacific oyster Crassostrea gigas, pearl oyster Pinctada fucata, freshwater snail Biomphalaria glabrata, owl limpet Lottia gigantea, California two-spot octopus Octopus bimaculoides, and California sea hare Aplysia californica) have been released (Zhang et al., 2012a; Takeuchi et al., 2012; Simakov et al., 2013; Albertin et al., 2015). These six molluscs, including four free-moving and two sessile species, can represent the end-lines of different and diverse evolutionary paths. The Pacific oyster and pearl oyster are representative bivalve molluscs. They are sessile filter-feeders with a bivalved shell. B. glabrata, L. gigantea and A. californica belong to the Gastropoda, which use the foot for crawling (Heller, 2015). Octopuses, a major group of the molluscan class Cephalopoda, have changed dramatically over the course of their evolution. They are active, resourceful predators with a rich behavioral repertoire (Albertin et al., 2015). The availability of molluscan genomes provides a unique opportunity to elucidate the adaptive evolution of the sessile molluscs at the genome level.

Here, we conducted genome-wide scans for the available molluscan genome sequences to identify the genes showing evidence of adaptive evolution along the sessile mollusc lineage. The findings will help elucidate the underlying molecular basis of adaptation to the sessile lifestyle.

# 2 Methods

The protein-coding sequences for five mollhefove mL.rn17u7(e(42 0 Td (, )Tj /TT-)JJ -0.4331 Tc 000215 Tw 2.O.2(o)- b6(p

hypergeometric test, and the calculated *P*-value was gone through FDR correction, taking FDR < 0.05 as a threshold.

## 3 Results and Discussion

Totally 303 single copy 1:1 orthologs from six available molluscan genomes were identified. Saturation of nucleotide substitution can bias evolutionary inferences, and result in an underestimation of d*S* and an inflation of dN/dS ( $\omega$ ). Therefore, we only retained 286 orthologs without significant substitution saturation for further




Fig.2 Distribution of  $\omega$  values obtained for all the genes on branch S using the branch model. Proportion of genes in the different  $\omega$  regions are indicated by the *Y*-axis with the  $\omega$  regions on the *X*-axis.

analysis. In order to estimate the impact of natural selection acting on genes, the one-rate model, which provides a single estimate of  $\omega$  across all codons and lineages, was utilized. All the 286 orthologs had  $\omega$  values in the range of 0.001–0.15, and no extreme  $\omega$  value was detected. The branch model (model = 2) was used to obtain the dN/dS values for all the genes on the labeled branch (branch S, Fig.1). The results showed that 61.19% of the genes had  $\omega$  values larger than 1 (Fig.2), suggesting a dramatic change in the selective pressures acting on the genes on branch S, thereby providing potential explanations for the emergence of new body plans, organs and lifestyles on branch S.

To detect signals of positive selection that may have occurred on branch S, the improved branch-site model was used in this study. This model can detect positive selection acting on a few sites on particular specified lineages. As a result, 40 genes showed significant evidence of positive selection (FDR < 0.2) on branch S. In order to understand the biological significance of the PSGs, we classified them according to the scheme established by the Gene Ontology project. Of the 40 PSGs, 22 PSGs were classified into 17 GO terms (Fig.3). The result revealed the most GO hits were in 'cellular component', followed by 'biological process' and 'molecular function'. Within the 'biological process' category, the most GO hits were in 'metabolic process'. Nine PSGs were related to catalytic activities within the molecular function category, including transferase activity, oxidoreductase activity, and hydrolase activity.




Fig.3 Histogram of the GO annotations of PSGs. The *X*-axis represents GO terms belonging to three categories and the *Y*-axis represents gene numbers of each term.

Among the 40 PSGs, 13 PSGs were assigned to ten pathways (Table 1). Enrichment analysis revealed that two pathways, Oxidative phosphorylation (OXPHOS) and TGF-beta signaling pathway, were significantly enriched among the PSGs. Surprisingly, genes NDUFB10, AT-PeF0B and Cyt1 were involved in Oxidative phosphorylation pathway under positive selection in *C. gigas* and *P. fucata*. This result was in strong contrast to most previous studies which detected association of positive selection on OXPHOS genes with evolution of a variety of energetically demanding characteristics (Shen *et al.*, 2010; Shen *et al.*, 2012). A common expectation is that OXPHOS functional efficiency is important to energy intensive processes, and positive selection on OXPHOS genes would lead to enhanced organismal performance, such as powered flight (Shen *et al.*, 2010). However, strong positive selection on OXPHOS genes detected among fishes with low aerobic performance did not match simple expectations, suggesting that the adaptation of OXPHOS involved many factors beyond enhancing ATP production

in high performance taxa, and would not be readily predicted based solely on organismal performance (Zhang and Broughton, 2015). In this study, the result showing positive selection on OXPHOS genes in sessile molluscs was consistent with those in low-performance fishes. The organismal fitness in sessile molluscs may be increased by a modified OXPHOS regulatory system or even a reduction in OXPHOS efficiency.

Pathway	Number of positively selected genes with pathway annotation	P-value	Q-value
Oxidative phosphorylation	3 (27.27%)	0.00095	0.00953
TGF-beta signaling pathway	2 (18.18%)	0.00822	0.04109
Pantothenate and CoA biosynthesis	1 (9.09%)	0.02841	0.09470
Nicotinate and nicotinamide metabolism	1 (9.09%)	0.07676	0.19190
Drug metabolism-other enzymes	1 (9.09%)	0.10318	0.20635
Ribosome biogenesis in eukaryotes	1 (9.09%)	0.15396	0.25660
Ribosome	1 (9.09%)	0.23989	0.31784
Pyrimidine metabolism	1 (9.09%)	0.25427	0.31784
Wnt signaling pathway	1 (9.09%)	0.34018	0.37798
Lysosome	1 (9.09%)	0.38514	0.38514

Table 1 Results of pathway enrichment analysis of PSGs from branch S

TGF-beta signaling plays a key role in the patterning of metazoan body plans and growth (Kenny et al., 2014). It is widely regarded as a 'module' capable of co-option into novel functions. In gastropods, TGF-beta signaling pathway has been reported as a major driving force of torsion by regulating asymmetric cell proliferation (Kurita and Wada, 2011). During the evolution of molluscs leading to bivalves, the most prominent morphological change was that a single dorsal shell was doubled (Kin et al., 2009). A member of TGF-beta family, *dpp*, was found to be very important for establishing the characteristic shape of the shell anlagen in Japanese spiny oyster, suggesting that TGF-beta signaling pathway played a pivotal role in the morphological transition to two shells in bivalve (Kin et al., 2009). In this study, two genes (BAMBI and ID1) in TGF-beta signaling pathway were found undergoing positive selection, which may be associated with the great changes in body plans and organs of the sessile molluscs.

A little unexpectedly, we did not find any known genes involved in immune and stress responses under positive selection in the two sessile bivalves, considering they may face tremendous exposure to microbial pathogens and inhabit in highly stressful and widely changing environments. This result is probably because only single copy 1:1 orthologs were selected to identify the natural selection in our study. Recent studies have revealed massive expansion of immune genes in sessile bivalves, suggesting that the expansion was also important for the evolution and adaptation of bivalve molluscs (Zhang *et al.*, 2012a; Guo *et al.*, 2015; Takeuchi *et al.*, 2016).

## 4 Conclusions

In summary, we conducted the first genome-wide scan for positively selected genes to investigate the genetic basis of attached lifestyle in the sessile molluscs, providing a new insight into the molecular landscape of sessile adaptation in molluscs. Although only six molluscan species were included in the analysis, it is clear that positive selection is indeed an important source of evolutionary innovation and plays a critical role in the evolution of the sessile molluscs. Along with the development of the genome sequencing techniques, assembly of higher-quality genomes, and more accurate gene annotation, the availability of more molluscan genomes will provide more resources for the investigation of adaption evolution in molluscs.

#### Acknowledgements

This study was supported by the grants from the National Natural Science Foundation of China (No. 3137 2524), the Shandong Seed Project, Shandong Province (No. 2014GHY115002), the Fundamental Research Funds for the Central Universities (No. 201762014), and Qingdao National Laboratory for Marine Science and Technology (No. 2016LMFS-A06).

### References

- Albertin, C. B., Simakov, O., Mitros, T., Wang, Z. Y., Pungor, J. R., Edsinger-Gonzalez, E., Brenner, S., Ragsdale, C. W., and Rokhsar, D. S., 2015. The octopus genome and the evolution of cephalopod neural and morphological novelties. *Nature*, 524: 220-224.
- Bayne, B. L., and Newell, R. C., 1983. Physiological energetics of marine molluscs. In: *The Mollusca, Vol.* 4. Saleuddin, A. S. M., and Wilbur, K. M., eds., Academic Press, New York, 407-515.
- Benjamini, Y., and Hochberg, Y., 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B–Statistical Methodology*, **57**: 289-300.
- Cao, X., Sun, Y. B., Irwin, D. M., Wang, G. D., and Zhang, Y. P., 2015. Nocturnal to diurnal transition in the common ancestor of haplorrhines: Evidence from genomic-scan for positively selected genes. *Journal of Genetics and Genomics*, 42: 33-37.
- Darriba, D., Taboada, G. L., Doallo, R., and Posada, D., 2011. ProtTest 3: Fast selection of best-fit models of protein evolution. *Bioinformatics*, 27: 1164-1165.
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology*, **59**: 307-321.
- Guo, X., He, Y., Zhang, L., Lelong, C., and Jouaux, A., 2015. Immune and stress responses in oysters with insights on ad-

aptation. Fish and Shellfish Immunology, 46: 107-119.

- Heller, J., 2015. *Sea Snails: A Natural History*. Springer, New York, 354pp.
- Kenny, N. J., Namigai, E. K. O., Dearden, P. K., Hui, J. H. L., Grande, C., and Shimeld, S. M., 2014. The Lophotrochozoan TGF-β signalling cassette-diversification and conservation in a key signalling pathway. *International Journal of Developmental Biology*, **58**: 533-549.
- Kin, K., Kakoi, S., and Wada, H., 2009. A novel role for *dpp* in the shaping of bivalve shells revealed in a conserved molluscan developmental program. *Developmental Biology*, **329**: 152-166.
- Kocot, K. M., Cannon, J. T., Todt, C., Citarella, M. R., Kohn, A. B., Meyer, A., Santos, S. R., Schander, C., Moroz, L. L., Lieb, B., and Halanych, K. M., 2011. Phylogenomics reveals deep molluscan relationships. *Nature*, **477**: 452-456.
- Kurita, Y., and Wada, H., 2011. Evidence that gastropod torsion is driven by asymmetric cell proliferation activated by TGF-β signaling. *Biology Letters*, **7**: 759-762.
- Meng, Y., Zhang, W., Zhou, J., Liu, M., Chen, J., Tian, S., Zhuo, M., Zhang, Y., Zhong, Y., Du, H., and Wang, X., 2015. Genome-wide analysis of positively selected genes in seasonal and non-seasonal breeding species. *PLoS One*, **10** (5): e012 6736.
- Nery, M. F., González, D. J., and Opazo, J. C., 2013. How to make a dolphin: Molecular signature of positive selection in cetacean genome. *PLoS One*, **8** (6): e65491.
- Ponder, W. F., and Lindberg, D. R., 2008. *Phylogeny and Evolution of the Mollusca*. University of California Press, Berkeley and Los Angeles, 469pp.
- Shen, Y. Y., Liang, L., Zhu, Z. H., Zhou, W. P., Irwin, D. M., and Zhang, Y. P., 2010. Adaptive evolution of energy metabolism genes and the origin of flight in bats. *Proceedings of the National Academy of Sciences*, **107**: 8666-8671.
- Shen, Y. Y., Zhou, W. P., Zhou, T. C., Zeng, Y. N., Li, G. M., Irwin, D. M., and Zhang, Y. P., 2012. Genome-wide scan for bats and dolphin to detect their genetic basis for new locomotive styles. *PLoS One*, 7 (11): e46455.
- Simakov, O., Marletaz, F., Cho, S. J., Edsinger-Gonzales, E., Havlak, P., Hellsten, U., Kuo, D. H., Larsson, T., Lv, J., Arendt, D., Savage, R., Osoegawa, K., de Jong, P., Grimwood, J., Chapman, J. A., Shapiro, H., Aerts, A., Otillar, R. P., Terry, A. Y., Boore, J. L., Grigoriev, I. V., Lindberg, D. R., Seaver, E. C., Weisblat, D. A., Putnam, N. H., and Rokhsar, D. S., 2013. Insights into bilaterian evolution from three spiralian genomes. *Nature*, 493: 526-531.
- Sun, Y. B., Zhou, W. P., Liu, H. Q., Irwin, D. M., Shen, Y. Y., and Zhang, Y. P., 2013. Genome-wide scans for candidate genes involved in the aquatic adaptation of dolphins. *Genome Biology and Evolution*, 5: 130-139.
- Takeuchi, T., Kawashima, T., Koyanagi, R., Gyoja, F., Tanaka, M., Ikuta, T., Shoguchi, E., Fujiwara, M., Shinzato, C., Hisata,

- K., Fujie, M., Usami, T., Nagai, K., Maeyama, K., Okamoto,
- K., Aoki, H., Ishikawa, H., Masaoka, T., Fujiwara, A., Endo, K., Endo, H., Nagasawa, H., Kinoshita, S., Asakawa, S., Watabe, S., and Satoh, N., 2012. Draft genome of the pearl oyster *Pinctada fucata*: A platform for understanding bivalve biology. *DNA Research*, **19** (2): 117-130.
- Takeuchi, T., Koyanagi, R., Gyoja, F., Kanda, M., Hisata, K.,
  - Fujie,earDN0arcZo.6(.ical 0.0627 Tw 21.-N0ar)8.be, S., andLet /Ts -0.0005