

Hybridization between “Haida No. 1” and Orange-shell line of the Pacific oyster reveals high heterosis in survival

Yuanxin Liang^a, Guohan Zhang^a, Gaowei Jiang^a, Yiming Hu^a, Jiafeng Fang^a, Yong Chi^a, Chengxun Xu^a, Weiguo Liu^c, Haijun Liu^c, Qi Li^{a,b,*}

^a Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, Qingdao 266003, China

^b Laboratory for Marine Fisheries Science and Food Production Processes, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China

^c Rushan Huaxin Foodstuffs Co., Ltd, Rushan 264509, China

ARTICLE INFO

Keywords:

Summer mass mortality
Heterosis
Crassostrea gigas
Inbred line
Selected line

ABSTRACT

In order to assess the heterosis of the hybrid between the selected lines and develop a new hybrid variety of the Pacific oyster (*Crassostrea gigas*) with excellent survival and growth characteristics, a diallel crosses between selected “Haida No.1” (H, 12th generation) line and Orange-shell (O, 9th generation) line of *C. gigas* were carried out. Heterosis for growth and survival of two parental lines (HH, H × H and OO, O × O) and their reciprocal hybrids (HO, H × O and OH, O × H) were systematically assessed at both larval and grow-out stages. At larval stage, the two reciprocal hybrids exhibited heterosis in both survival and growth. Compared with HH, the survival rate of OH increased by nearly 23.00% at larval stage. Moreover, the heterosis in survival (average: 35.27%) was higher than that in growth (average: 8.03%) at larval stage. During grow-out phase, reciprocal hybrids showed excellent growth and survival traits at three commercial culturing areas. The high-parent heterosis for survival of hybrids at Rushan were higher than those at Rongcheng or Huangdao. Notably, the survival rate of OH increased by nearly 32.00% compared with HH at Rushan. Among three sites, the high-parent heterosis in survival varied from 31.18% to 141.18% at day 450. Overall, OH (Orange-shell line × “Haida No. 1” line) exhibited superior growth and survival traits among four crosses, which could be prioritized as a new alternative variety for commercial oyster cultivation.

1. Introduction

The Pacific oyster (*C. gigas*) is one of the most economically important aquatic species in the world. In recent years, genetic improvement for economic traits of *C. gigas* have been widely conducted, such as disease tolerance (Divilov et al., 2021), growth (de Melo et al., 2016), yield (Langdon et al., 2003; Rawson and Feindel, 2012), and survival (Dégremont et al., 2010). Nonetheless, several problems remain, e.g., summer mass mortality of oysters (SMM). SMM have caused serious economic losses and have severely hampered the development of the oyster industry (Du et al., 2021; Pernet et al., 2012; Evans and Langdon, 2006; Azéma et al., 2015; Pernet et al., 2010). In France, 7-month-old oysters reared in Marennes-Oléron Bay suffered 50% mortality during 17 days in August 2009 (Dégremont, 2011). In northern China, the mortality rate of *C. gigas* was more than 60% in summer months (Yang et al., 2021). Moreover, SMM breaks out in oysters, regardless of ploidy levels and origins (Pernet et al., 2010; Azéma et al., 2016).

Crossbreeding combining favorable characteristics of both parents has been extensively applied in genetic breeding of organisms. In aquaculture, hybridization was commonly used to improve flesh quality, increase growth and survival rates, enhance disease and stress tolerance, as well as manipulate sex ratios, produce sterile animals (Bartley et al., 2001; Dang et al., 2011; Liang et al., 2014; Zhang et al., 2017a; Liang et al., 2018; Tan et al., 2020). For example, hybrids derived from two stocks of *C. hongkongensis* exhibited superior heterosis in survival and living weight (Zhang et al., 2017b). The survival and whole weight of *C. sikamea* could be significantly improved through crossbreeding between Chinese and American *C. sikamea* populations (Ma et al., 2022). Two hybrid strains with superior growth and survival could be obtained by hybridization between 4 geographical subpopulations of sea scallops (Wang and Côté, 2012). Additive genetic variation in yield traits has been accumulated by repeated selection. Non-additive genetic variation can be utilized in the offspring through crossbreeding. Therefore, hybridization between selected lines offer the possibility of exploiting both

* Corresponding author.

E-mail address: qili66@ouc.edu.cn (Q. Li).

<https://doi.org/10.1016/j.aquaculture.2022.737945>

Received 2 December 2021; Received in revised form 10 January 2022; Accepted 16 January 2022

Available online 21 January 2022

0044-8486/© 2022 Elsevier B.V. All rights reserved.

additive genetic variation accumulated within lines and non-additive genetic variation between lines (Han et al., 2020; Hedgecock and Davis, 2007). High heterosis in phenotypic characteristics was observed in crosses between selection lines of *Ruditapes philippinarum* (Zhang et al., 2014) and *C. gigas* (Kong et al., 2017; Han et al., 2020). Besides, the high-yield offspring was

To determine the effects of the genotypes (hybrid groups vs. purebred groups) and environmental factors (Rushan vs. Rongcheng vs. Huangdao) on the survival and growth of *C. gigas* during juvenile and adult stage. A two-factor ANOVA was employed as the following model (Mallet and Haley, 1983; Cruz and Ibarra, 1997):

$$Y_{ijk} = \mu + G_i + E_j + (G \times E)_{ij} + e_{ijk}$$

Where Y_{ijk} indicates the mean phenotypic value (shell height, survival rate, etc) of the k replicate from the type i group in site j . G_i is the genotype effect on phenotypic value (shell height, survival rate, etc) ($i = 1, 2$), E_j is the environmental effect on the phenotypic value (shell height, survival rate, etc) ($j = 1, 2, 3$), $(G \times E)_{ij}$ represents the interaction effect of the group type and site, and e_{ijk} is the random observation error ($k = 1, 2, 3$).

3. Results

3.1. Growth, survival and heterosis at larval stage

No significant difference was observed in shell height between two reciprocal hybrids (Fig. 1A). The shell height of OO was lower than those of the other three groups at larval stage. From day 15, two reciprocal hybrids were significantly larger than two purebred lines in shell height ($P < 0.05$), with the order of $OH > HO > HH > OO$. On day 20, average shell height of HO ($285.67 \pm 35.87 \mu\text{m}$) was slightly larger than that of OH ($275.25 \pm 24.75 \mu\text{m}$). The shell heights of two hybrid groups were significantly larger than those of two purebred lines at day 20 ($P < 0.05$),

with a high-parent heterosis $H_{(HO/HH)}$ of 15.51% and $H_{(OH/HH)}$ of 11.30% (Table 1).

On day 15, the mean survival rates of the hybrid groups were significantly higher than those of purebred groups ($P < 0.05$) (Fig. 1B). However, no significant difference was observed between the two hybrids. The OH group exhibited highest survival rate among the four groups on day 20, with a heterosis of 46.00% compared to HH (Table 1). Notably, the survival rate of OO group ($37.66 \pm 3.06\%$) was significantly lower than those of the other three groups ($P < 0.05$). Moreover, heterosis for survival in the hybrid groups (2.84–46.00%) gradually increased with larval growth (Table 1).

3.2. Growth, survival and heterosis at juvenile stage

Two-factor ANOVA showed that growth and survival traits for four experimental groups were significantly influenced by genotypes and environments (Table 2). The shell heights of four groups in Rushan were larger than those in Rongcheng and Huangdao (Fig. 2). Two hybrid groups exhibited heterosis in shell height compared to HH, with mid-parent heterosis ranged from 0.52% to 18.33% at three culturing areas. At day 450, significant difference in shell height was observed among four groups in Rongcheng ($P < 0.05$). Besides, the shell heights of two hybrid groups were larger than that of HH, with high-parent heterosis of 7.74% and 15.41% for HO and OH, respectively (Table 3). Moreover, the shell heights of OO were significantly smaller than those of the other three groups at three areas ($P < 0.05$), followed the order of $OH > HO > HH > OO$.

At day 450, the living weights of hybrid groups were significantly ($P < 0.05$) higher than those of purebred groups at Rongcheng (Fig. 2). Moreover, OH was significantly ($P < 0.05$) heavier than the other three groups, with values of high-parent heterosis at 18.44% and 22.00% in Rushan and Huangdao, respectively. Besides, the heterosis for living weight was higher at Rongcheng than at Rushan or Huangdao. While the living weights of four experimental groups at Rushan were highest among three environments (Table 4).

The mean survival rates of two hybrid crosses were higher than those of the two purebred groups at three environments (Fig. 2). At day 90, two hybrid lines in Rushan showed negative high-parent heterosis, with values at -3.07% for HO and -1.15% for OH (Table 5). Similar results were also observed in Rongcheng and Huangdao. However, the survival rates of hybrid crosses (HO: 86.66%, OH: 85.66%) were higher than those of their parental groups (HH: 73.33%, OO: 64.00%) on day 180 in Rongcheng. Moreover, the survival rate of OO were significantly ($P < 0.05$) lower than those of the other three groups, except on day 90 at three environments. Notably, on day 450, which indicates oysters had tolerated the summer for the second time, the survival rates of hybrid crosses were significantly ($P < 0.05$) higher than purebred groups at three culture areas. Meanwhile, OH group exhibited higher heterosis in survival than HO group, with high-parent heterosis ranged from 102.15% to 141.18% at three environments. Besides, the heterosis for




Fig. 1. Shell height and survival rate for two purebred groups (HH and OO) and two hybrid groups (HO and OH) at larval stage. A: the shell height of four groups; B: the survival rate of four groups. H and O indicate “Haida No. 1” line and Orange-shell line of the Pacific oyster, respectively. Two purebred groups: HH ($H \times H$), OO ($O \times O$); and their reciprocal hybrids: HO ($H \times O$), OH ($O \times H$). Different superscript letters in the same day indicate significant difference ($P < 0.05$) among four groups.

Table 1

Heterosis (M and H) for survival rate and shell height of “Haida No. 1” (H) line, Orange-shell (O) line and their reciprocal hybrids (HO and OH) at larval stage.

| Items | Heterosis (%) | Day 1 | Day 5 | Day 10 | Day 15 | Day 20 |
|---------------|---------------|-------|-------|--------|--------|--------|
| Survival rate | M_{F1} | – | 5.25 | 17.98 | 54.35 | 63.50 |
| | $H_{(HO/HH)}$ | – | 2.84 | 18.72 | 29.02 | 40.67 |
| | $H_{(OH/HH)}$ | – | 3.19 | 21.00 | 28.50 | 46.00 |
| Shell height | M_{F1} | 6.07 | 5.54 | 1.89 | 10.75 | 15.87 |
| | $H_{(HO/HH)}$ | 4.47 | 4.62 | 3.06 | 8.87 | 15.51 |
| | $H_{(OH/HH)}$ | 4.50 | 3.11 | 1.139 | 9.59 | 11.30 |

H and O indicate “Haida No. 1” and Orange-shell lines of the Pacific oyster, respectively. Two purebred groups: HH ($H \times H$), OO ($O \times O$); and two reciprocal hybrids: HO ($H \times O$), OH ($O \times H$). M_{F1} represents the mid-parent heterosis of the two hybrids; $H_{(HO/HH)}$ and $H_{(OH/HH)}$ means the high-parent heterosis of HO and OH, respectively.

Table 2

Two-factor analysis of variance (ANOVA) showing the genotype (G) and environment factor (E) effects for survival and growth of each experimental group at juvenile and adult stage.

| Items | Source | df | Survival rate | | Shell height | | Living weight | |
|---------|--------|----|---------------|-----------|--------------|-----------|---------------|-----------|
| | | | MS | F-value | MS | F-value | MS | F-value |
| Day 90 | E | 2 | 0.154 | <0.001*** | 0.002 | 0.747 | 2.319 | <0.001*** |
| | G | 1 | 0.002 | 0.426 | 0.775 | <0.001*** | 2.438 | <0.001*** |
| | G * E | 2 | 0.007 | 0.146 | 0.049 | 0.001** | 0.037 | 0.231 |
| Day 180 | E | 1 | 0.035 | 0.020** | 0.057 | <0.001*** | 1.220 | <0.001*** |
| | G | 2 | 0.650 | <0.001*** | 0.682 | <0.001*** | 2.784 | <0.001*** |
| | G * E | 2 | 0.001 | 0.913 | 0.012 | 0.101 | 0.056 | 0.035* |
| Day 270 | E | 2 | 0.032 | 0.003** | 0.171 | <0.001*** | 1.795 | <0.001*** |
| | G | 1 | 0.673 | <0.001*** | 0.444 | <0.001*** | 2.936 | <0.001*** |
| | G * E | 2 | 0.002 | 0.691 | 0.000 | 0.932 | 0.067 | 0.006** |
| Day 360 | E | 2 | 0.052 | 0.003** | 0.135 | <0.001*** | 1.565 | <0.001*** |
| | G | 1 | 0.628 | <0.001*** | 0.027 | 0.003** | 0.014 | 0.237 |
| | G * E | 2 | 0.001 | 0.903 | 0.226 | <0.001*** | 1.209 | <0.001*** |
| Day 450 | E | 1 | 0.727 | 0.196 | 0.080 | <0.001*** | 1.134 | <0.001*** |
| | G | 2 | <0.001 | <0.001*** | 0.476 | <0.001*** | 1.855 | <0.001*** |
| | G * E | 2 | 0.011 | 0.981 | 0.017 | 0.002** | 0.083 | <0.001*** |

* indicates $P < 0.05$; ** indicates $P < 0.01$; *** indicates $P < 0.001$.

survival of hybrid groups were highest at Rushan.

3.3. Shell color of the hybrid cohorts

The 1000 hybrids from Rushan were classified by eye observation based on shell pigmentation of the left valve. Hybrid cohorts had two types on color patterns. The shell pattern of 58.4% of the progenies is the same as that of "Haida No. 1" line with or without radial stripes on left shell, while 41.6% of the hybrids exhibited purple colors (ranging from pale to dark purple) on both shells (Fig. 3).

4. Discussion

Improving yield traits such as growth rate and survival rate has been the main objective of genetic breeding for aquatic animals. Here, we compared the phenotypic values of the Pacific oyster "Haida No. 1" line, the Orange-shell line and their reciprocal crosses. The results showed that the four groups of oysters differed significantly in growth and survival at three environments.

4.1. Growth

The results of the study revealed that hybrid crosses exhibited heterosis in shell height, with values of mid-parent heterosis ranging from 1.89% to 15.87% at larval stage and 13.46% to 24.48% at grow-out stage. The positive association between growth and heterozygosity in marine mollusks is a common consequence of crossbreeding (Zouros and Mallet, 1989; Sheridan, 1981). Besides, the successful use of heterosis is attributed to an increase in heterozygosity (Mittton and Grant, 1984). Therefore, growth difference in this study may be interpreted as higher heterozygosity in the hybrids than in purebreds. In this study, hybrid crosses exhibited low heterosis in growth, and the similar results were also demonstrated in *C. gigas* (Kong et al., 2017; Han et al., 2020) and *Argopecten irradians* (Zhang et al., 2007; Wang and Li, 2010). However, high heterosis in shell height and living weight was found in reciprocal hybrids derived from *C. hongkongensis* and *C. sikamea* (Zhang et al., 2017a). The growth difference in hybrids may be caused by genetic differences between parents (Sheridan, 1997) and environmental conditions (Evans and Langdon, 2006). In addition, different degrees of inbreeding among parental populations may be another important cause of the above phenomenon (Rawson and Feindel, 2012). Moreover, the shell height and living weight were higher in HH than in OO. However, the heterosis for growth-related traits in OH were higher than that in HO. This phenomenon indicates the non-additive effects were critical for oyster growth (Hedgecock et al., 1995; Hedgecock and Davis, 2007).

4.2. Survival

In this study, the hybrid groups exhibited higher survival rate than purebred groups at both larval and grow-out stage. Especially, OH showed a significant survival advantage over the other three groups, with high-parent heterosis of 141.18% at Rushan, which was probably due to better adaptation of reciprocal hybrids than their parental counterparts under certain environmental conditions (Zhang et al., 2017b). On the other hand, heterozygosity and hybridization may have masked the deleterious or even lethal recessive survival-related genes in the parental lines (Wang and Li, 2010; Burke and Arnold, 2001). Moreover, "Haida No. 1" line originated from cultured stocks in Rushan, while the Orange-shell line was established based on only four *C. gigas*. Thus, genetic variation may exist between these two parental lines. The magnitude of heterosis between two particular lines depends on the square of the difference in gene frequency between crossed parents. Moreover, if no different gene frequency was existed between the parents crossed, there will be no heterosis. Nevertheless, the heterosis will be greatest when one particular trait-related loci was fixed in one parent and the other loci was fixed in the other parent (Falconer, 1981). Besides, high heterosis in survival suggested crossbreeding between selected lines is an efficient method to improve the survival of *C. gigas*.

The heterosis in survival was higher than that in growth throughout the life stage. This may be attributed to the fact that the two parental populations have been selected for generations with directional selection for growth rather than for survival. Thus, the survival-related loci of the two parents may be more diverse and complementary than growth-related loci (Han et al., 2020). On the other hand, fitness-related traits (survival rate, etc) are more likely to exhibit directional dominance compared to morphological-related traits (shell height, etc) (Lynch and Walsh, 1998), and the magnitude of heterosis depends largely on directional dominance of genes involved in target traits (Falconer and Mackay, 1996). Thus, the heterosis between two reciprocal groups is greater for survival than for growth.

4.3. Effects of generations and inbreed lines

The two reciprocal hybrids exhibited extraordinary performances in terms of survival and growth. Superior performance of hybrids could be partially explained by the combination of selection and crossbreeding. The Orange-shell line has been successively selected for 9 generations. Meanwhile, the "Haida No. 1" line has been successively selected for 12 generations with fast growth. Theoretically, when the parental populations have been selected for generations, the hybrid progeny will exhibit high heterosis due to the accumulation of plenty of different non-




Fig. 2. Shell height, living weight and survival rate for two purebred groups (HH and OO) and two hybrid groups (HO and OH) among three sites. A: the shell height of four groups at Rongcheng; B: the living weight of four groups at Rongcheng; C: the survival rate of four groups at Rongcheng; D: the shell height of four groups at Rushan; E: the living weight of four groups at Rushan; F: the survival rate of four groups at Rushan; G: the shell height of four groups at Huangdao; H: the living weight of four groups at Huangdao; I: the survival rate of four groups at Huangdao. Different superscript letters in the same day indicate significant difference ($P < 0.05$) among four groups.

Table 3

Heterosis (*M* and *H*) for shell height in “Haida No. 1” (H) line, Orange-shell (O) line and their reciprocal hybrids (HO and OH) at grow-out stage in three culturing environments.

| Heterosis (%) | Day 90 | Day 180 | Day 270 | Day 360 | Day 450 |
|-----------------------------|--------|---------|---------|---------|---------|
| Rongcheng | | | | | |
| <i>M</i> _{F1} | 10.91 | 20.09 | 15.41 | 19.66 | 24.48 |
| <i>H</i> _(HO/HH) | 4.53 | 4.89 | 3.03 | 6.61 | 7.74 |
| <i>H</i> _(OH/HH) | 0.63 | 2.62 | 0.52 | 11.41 | 15.41 |
| Rushan | | | | | |
| <i>M</i> _{F1} | 34.43 | 26.23 | 17.85 | 15.82 | 13.46 |
| <i>H</i> _(HO/HH) | 18.33 | 14.71 | 7.24 | 6.33 | 6.32 |
| <i>H</i> _(OH/HH) | 17.90 | 13.57 | 12.44 | 9.92 | 8.10 |
| Huangdao | | | | | |
| <i>M</i> _{F1} | 24.34 | 15.54 | 16.64 | 16.89 | 15.01 |
| <i>H</i> _(HO/HH) | 4.33 | 1.13 | 3.12 | 4.28 | 1.54 |
| <i>H</i> _(OH/HH) | 9.27 | 2.51 | 4.95 | 8.29 | 8.31 |

Table 4

Heterosis (*M* and *H*) for living weight in “Haida No. 1” (H) line, Orange-shell (O) line and their reciprocal hybrids (HO and OH) at grow-out stage among three culturing environments.

| Heterosis (%) | Day 90 | Day 180 | Day 270 | Day 360 | Day 450 |
|-----------------------------|--------|---------|---------|---------|---------|
| Rongcheng | | | | | |
| <i>M</i> _{F1} | 38.62 | 56.87 | 60.67 | 57.69 | 53.85 |
| <i>H</i> _(HO/HH) | 19.52 | 13.74 | 28.11 | 27.60 | 27.76 |
| <i>H</i> _(OH/HH) | 47.89 | 73.78 | 73.46 | 69.93 | 66.38 |
| Rushan | | | | | |
| <i>M</i> _{F1} | 48.61 | 44.03 | 35.95 | 29.06 | 24.18 |
| <i>H</i> _(HO/HH) | 6.96 | 10.51 | 9.80 | 6.49 | 3.87 |
| <i>H</i> _(OH/HH) | 16.57 | 25.49 | 22.35 | 20.46 | 18.44 |
| Huangdao | | | | | |
| <i>M</i> _{F1} | 29.98 | 31.74 | 44.71 | 37.03 | 34.89 |
| <i>H</i> _(HO/HH) | -5.49 | 0.18 | 7.38 | 4.76 | 4.19 |
| <i>H</i> _(OH/HH) | -4.10 | 8.12 | 26.77 | 23.11 | 22.00 |

Table 5

Heterosis (*M* and *H*) for survival in “Haida No. 1” (H) line, Orange-shell (O) line and their reciprocal hybrids (HO and OH) at grow-out stage in three culturing environments.

| Heterosis (%) | Day 90 | Day 180 | Day 270 | Day 360 | Day 450 |
|-----------------------------|--------|---------|---------|---------|---------|
| Rongcheng | | | | | |
| <i>M</i> _{F1} | 0.00 | 25.49 | 47.47 | 57.98 | 105.30 |
| <i>H</i> _(HO/HH) | 0.00 | 18.18 | 27.71 | 27.91 | 31.18 |
| <i>H</i> _(OH/HH) | -1.07 | 16.82 | 36.14 | 63.57 | 102.15 |
| Rushan | | | | | |
| <i>M</i> _{F1} | 5.36 | 29.00 | 50.79 | 77.33 | 138.60 |
| <i>H</i> _(HO/HH) | -3.07 | 7.08 | 25.00 | 27.84 | 58.82 |
| <i>H</i> _(OH/HH) | -1.15 | 17.45 | 33.78 | 86.60 | 141.18 |
| Huangdao | | | | | |
| <i>M</i> _{F1} | -1.37 | 30.16 | 46.59 | 69.42 | 133.62 |
| <i>H</i> _(HO/HH) | -3.86 | 12.04 | 23.57 | 48.08 | 47.22 |
| <i>H</i> _(OH/HH) | -1.93 | 15.74 | 36.94 | 87.50 | 129.17 |

additive genetic variation, and the more successive generations of the parents, the higher heterosis in the hybrids (Sheridan, 1997).

The orange shell line used in this study is an inbred line established based only on four *C. gigas*. When this inbred orange-shell line was crossed with “Haida No. 1” line as one parent, the hybrid groups exhibited favorable performances, especially in survival. This phenomenon probably due to the yield-related loci of these two parents may be more complementary (Han et al., 2020). Meanwhile, this result indicates inbred lines have great application potential in aquaculture (Hedgecock et al., 1995; Hedgecock and Davis, 2007). Nevertheless, inbreeding and crossing alone cannot produce any improvement unless the inbred lines derived from different foundation populations. Moreover, parental populations must be selected at some stage if any improvement is to be made (Gjerde, 1988).

4.4. Difference performances existed in different crosses and environments

The heterosis for survival of OH was greater than that of HO throughout the life stage. This indicated the importance of evaluating the performances of two reciprocal hybrids before choosing the paternal and maternal line for commercial seed production (Zhang et al., 2007). Asymmetric performance between reciprocal crosses were also found in the Kumamoto oyster *C. sikamea* (Ma et al., 2022), the Hongkong oyster *C. hongkongensis* (Zhang et al., 2017a), the bay scallop *A. irradians* (Zhang et al., 2007) and the boring giant clam *Tridacna crocea* (Zhang et al., 2020). The performance variation between reciprocal hybrids were probably due to sex-linked genes, cytoplasmic inheritance, parental effects and extra-nuclear effects (Kong et al., 2017; Zhang et al., 2017a).

Both survival and growth traits of four groups varied among all three culturing environments. Notably, greater high-parent heterosis for survival and growth were found at Rushan. This difference may be caused by environmental condition, such as water circulation, salinity, temperature and food (Xu et al., 2019). Different performances among culture areas indicates the importance of evaluating the phenotypic traits of oysters among different environments to obtain greatest heterosis.

4.5. Application prospect of shell color variants

Aquatic organisms with special colors are not only of high scientific value, but also more favored by consumers in the market (Clydesdale, 1993; Alfnes et al., 2006). In this study, oysters with purple on both the left and right shells were unexpectedly found from the hybrid offspring. Therefore, it is necessary to employ them as parents to develop a new line with stable inheritance of purple shell color. The total left-shell pigmentation in *C. gigas* was under a high degree of genetic control and strongly influenced by additive genetic variation (Brake et al., 2004; Evans et al., 2009). Besides, the heritability estimates in growth-related traits of the Pacific oyster were medium to high (Evans and Langdon, 2006; Kong et al., 2015; Chi et al., 2021; Li et al., 2011). The relatively high heritability of traits recorded in oysters also indicated that artificial selection has a significant potential to improve the performance of this species (Kong et al., 2015; Zhang et al., 2019). Thus, selective breeding including family and mass selection could be prioritized for genetic improvement of purple-variant.

5. Conclusions

In summary, the growth and survival of “Haida No. 1” line, Orange-shell line and their reciprocal hybrids were systematically assessed under three commercial conditions. The reciprocal cross exhibited significant heterosis in terms of survival at both larval and grow-out stages. The high-parent heterosis of OH was 46.00% in survival at larval stage. Meanwhile, the survival rate of OH increased by nearly 32.00% compared with HH on day 450. Moreover, the heterosis for shell height and living weight of OH was higher than that of HO. Therefore, the results demonstrated that OH (Orange-shell line × “Haida No. 1” line) has significant potential for application in the oyster aquaculture.

Credit author statement

Yuanxin Liang: Methodology, Software, Writing-original draft. **Guohan Zhang:** Data curation, Methodology. **Gaowei Jiang:** Writing-review. **Yiming Hu:** Writing-review. **Jiafeng Fang:** Writing-review. **Yong Chi:** Data curation, Software. **Qi Li:** Conceptualization, Formal analysis, Methodology, Writing-review. **Chengxun Xu:** Supervision. **Weiguo Liu:** Software. **Haijun Liu:** Software.

Declaration of Competing Interest


The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by the grants from National Natural Science Foundation of China (31972789), the China Agriculture Research System Project (CARS-49), Earmarked Fund for Agriculture Seed Improvement Project of Shandong Province (2020LZGC016), and Industrial Development Project of Qingdao City (20-3-4-16-nsh).

Cruz, P., Ibarra, A.M., 1997. Larval growth and survival of two catarina scallop (*Argopecten circularis*, Sowerby, 1835) populations and their reciprocal crosses. *J. Exp. Mar. Biol. Ecol.* 212, 95–110. [https://doi.org/10.1016/S0022-0981\(96\)02742-6](https://doi.org/10.1016/S0022-0981(96)02742-6).

Dang, V.T., Speck, P., Doroudi, M., Smith, B., Benkendorff, K., 2011. Variation in the antiviral and antibacterial activity of abalone *Haliotis laevigata*, *H. rubra* and their in <https://doi.org/10.1016/j.aquaculture.2011.05.010>.



- Han, Z.Q., Li, Q., 2020. Mendelian inheritance of orange shell color in the Pacific oyster *Crassostrea gigas*. *Aquaculture* 516, 734616. <https://doi.org/10.1016/j.aquaculture.2019.734616>.
- Han, Z.Q., Li, Q., Liu, S.K., Yu, H., Kong, L.F., 2019. Genetic variability of an orange-shell line of the Pacific oyster *Crassostrea gigas* during artificial selection inferred from microsatellites and mitochondrial COI sequences. *Aquaculture* 508, 159–166. <https://doi.org/10.1016/j.aquaculture.2019.04.074>.
- Han, Z.Q., Li, Q., Liu, S.K., Kong, L.F., 2020. Crossbreeding of three different shell color lines in the Pacific oyster reveals high heterosis for survival but low heterosis for growth. *Aquaculture* 529, 735621. <https://doi.org/10.1016/j.aquaculture.2020.735621>.
- Hedgecock, D., Davis, J.P., 2007. Heterosis for yield and crossbreeding of the Pacific oyster *Crassostrea gigas*. *Aquaculture* 272, S17–S29. <https://doi.org/10.1016/j.aquaculture.2007.07.226>.
- Hedgecock, D., McGoldrick, D.J., Bayne, B.L., 1995. Hybrid vigor in Pacific oysters: an experimental approach using crosses among inbred lines. *Aquaculture* 137, 285–298. [https://doi.org/10.1016/0044-8486\(95\)01105-6](https://doi.org/10.1016/0044-8486(95)01105-6).
- Jiang, G.W., Li, Q., Xu, C.X., Liu, S.K., Kong, L.F., Yu, H., 2021. Reciprocal hybrids derived from *Crassostrea gigas* and *C. angulata* exhibit high heterosis in growth, survival and thermotolerance in northern China. *Aquaculture* 545, 737173. <https://doi.org/10.1016/j.aquaculture.2021.737173>.
- Kong, N., Li, Q., Yu, H., Kong, L.F., 2015. Heritability estimates for growth-related traits in the Pacific oyster (*Crassostrea gigas*) using a molecular pedigree. *Aquac. Res.* 46, 499–508. <https://doi.org/10.1111/are.12205>.
- Kong, L.F., Song, S.L., Li, Q., 2017. The effect of interstrain hybridization on the production performance in the Pacific oyster *Crassostrea gigas*. *Aquaculture* 472, 44–49. <https://doi.org/10.1016/j.aquaculture.2016.07.018>.
- Langeron, C., Evans, F., Jacobson, D., Blouin, M., 2003. Yields of cultured Pacific oysters *Crassostrea gigas* Thunberg improved after one generation of selection. *Aquaculture* 220, 227–244. [https://doi.org/10.1016/S0044-8486\(02\)00621-X](https://doi.org/10.1016/S0044-8486(02)00621-X).
- Li, Q., Wang, Q.Z., Liu, S.K., Kong, L.F., 2011. Selection response and realized heritability for growth in three stocks of the Pacific oyster *Crassostrea gigas*. *Fish. Sci.* 77, 643–648. <https://doi.org/10.1007/s12562-011-0369-0>.
- Liang, S., Luo, X., You, W.W., Luo, L.Z., Ke, C.H., 2014. The role of hybridization in improving the immune response and thermal tolerance of abalone. *Fish Shellfish Immunol.* 39, 69–77. <https://doi.org/10.1016/j.fsi.2014.04.014>.
- Liang, S., Luo, X., You, W.W., Ke, C.H., 2018. Hybridization improved bacteria resistance in abalone: evidence from physiological and molecular responses. *Fish Shellfish Immunol.* 72, 679–689. <https://doi.org/10.1016/j.fsi.2017.11.009>.
- Lynch, M., Walsh, B., 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer Associates Inc., Sunderland, Massachusetts.
- Ma, H.T., Lv, W.G., Qin, Y.P., Li, J., Li, X.Y., Liao, Q.L., Yu, Z.N., 2022. Aquaculture potential of two Kumamoto oyster (*Crassostrea sikamea*) populations and their reciprocal hybrids in southern China. *Aquaculture* 546, 737301. <https://doi.org/10.1016/j.aquaculture.2021.737301>.
- Mallet, A.L., Haley, L.E., 1983. Growth rate and survival in pure population matings and crosses of the oyster *Crassostrea virginica*. *Can. J. Fish. Aquat. Sci.* 40, 948–954. <https://doi.org/10.1139/f83-121>.
- Mitton, J.B., Grant, M.C., 1984. Associations among protein heterozygosity, growth rate, and developmental homeostasis. *Annu. Rev. Ecol. Syst.* 15, 479–499. <https://doi.org/10.1146/annurev.es.15.110184.002403>.
- Pernet, F., Barret, J., Marty, C., Moal, J., Le Gall, P., Boudry, P., 2010. Environmental anomalies, energetic reserves and fatty acid modifications in oysters coincide with an exceptional mortality event. *Mar. Ecol. Prog. Ser.* 401, 129–146. <https://doi.org/10.3354/meps08407>.
- Pernet, F., Barret, J., Le Gall, P., Corporeau, C., Dégremont, L., Lagarde, F., Keck, N., 2012. Mass mortalities of Pacific oysters *Crassostrea gigas* reflect infectious diseases and vary with farming practices in the Mediterranean Thau lagoon, France. *Aquac. Environ. Interact.* 2, 215–237. <https://doi.org/10.3354/aei00041>.
- Rawson, P., Feindel, S., 2012. Growth and survival for genetically improved lines of eastern oysters (*Crassostrea virginica*) and interline hybrids in Maine, USA. *Aquaculture* 326–329, 61–67. <https://doi.org/10.1016/j.aquaculture.2011.11.030>.
- Sheridan, J.P., 2001. Crossbreeding and heterosis. *Anim. Breed. Abstr.* 49, 131–144.
- Shey

improvement

of